A Appendix

The problem in Equation (2) is a convex optimization problem. It can be rewritten as

$$\underset{\alpha}{\text{minimize}} \max\left\{\frac{a_1}{b_1+\alpha}, \cdots, \frac{a_n}{b_n+\alpha}\right\} + \lambda\alpha, \quad \text{subject to} \quad \alpha > 0.$$
 (A.1)

Equation (A.1) is a convex optimization problem because the first term of the objective function is the sum of a pointwise maximum of convex functions, $a_i/(b_i + \alpha)$, which is convex in α .

Theorem 1. The sequence $\{f(\mathbf{x}^j)\}_{j=1,2,\dots}$ provided by Equation (7) converges to the local maximum of the density field.

Proof. $f(\mathbf{x})$ shown in Equation (3) is a bounded function because it is the sum of finite bounded kernel density functions. To prove the theorem, it is sufficient to show that the sequence $\{f(\mathbf{x}^j)\}_{j=1,2,\cdots}$ is strictly monotonically increasing, i.e., $f(\mathbf{x}^j) < f(\mathbf{x}^{j+1})$, if $\mathbf{x}^j \neq \mathbf{x}^{j+1}$.

From Equation (3),

$$f(\mathbf{x}^{j+1}) - f(\mathbf{x}^{j}) = \frac{c}{N} \sum_{i=1}^{N} \frac{1}{h_i} \left(k \left(\frac{\|\mathbf{d}(\mathbf{l}_i, \mathbf{x}^{j+1})\|^2}{h_i^2} \right) - k \left(\frac{\|\mathbf{d}(\mathbf{l}_i, \mathbf{x}^{j})\|^2}{h_i^2} \right) \right).$$
(A.2)

If a function, $\phi(z)$, is convex, the following inequality holds:

$$\phi(z_2) - \phi(z_1) \ge \phi'(z_1)(z_2 - z_1), \tag{A.3}$$

where ϕ' is the derivative of ϕ .

The profile, k(z), of the Gaussian kernel density function is convex and it satisfies Equation (A.3):

$$k(z_2) - k(z_1) \geq k'(z_1)(z_2 - z_1).$$
 (A.4)

 $\|\mathbf{z}\|^2$ is a convex function in \mathbf{z} where \mathbf{z} is a vector and thus,

$$\|\mathbf{z}_{2}\|^{2} - \|\mathbf{z}_{1}\|^{2} \ge 2\mathbf{z}^{\mathsf{T}} (\mathbf{z}_{2} - \mathbf{z}_{1}).$$
(A.5)

The perspective distance vector function, $d(l_i, z)$, is also convex in z because it is a linear-fractional function [1,2] which perserves the convexity and thus,

$$\mathbf{d}(\mathbf{l}_i, \mathbf{z}_2) - \mathbf{d}(\mathbf{l}_i, \mathbf{z}_1) \ge \nabla_{\mathbf{z}}(\mathbf{d}(\mathbf{l}_i, \mathbf{z}_1)) \left(\mathbf{z}_2 - \mathbf{z}_1\right).$$
(A.6)

From Equation (A.5) and (A.6), the following inequality holds:

$$\begin{aligned} \left\| \mathbf{d}(\mathbf{l}_{i},\mathbf{x}^{j+1}) \right\|^{2} &= \left\| \mathbf{d}(\mathbf{l}_{i},\mathbf{x}^{j}) \right\|^{2} &\geq 2\mathbf{d}(\mathbf{l}_{i},\mathbf{x}^{j})^{\mathsf{T}} \left(\mathbf{d}(\mathbf{l}_{i},\mathbf{x}^{j+1}) - \mathbf{d}(\mathbf{l}_{i},\mathbf{x}^{j}) \right) \\ &\geq 2\mathbf{d}(\mathbf{l}_{i},\mathbf{x}^{j})^{\mathsf{T}} \left(\left(\nabla_{\mathbf{x}} \mathbf{d}(\mathbf{l}_{i},\mathbf{x}^{j}) \right) \left(\mathbf{x}^{j+1} - \mathbf{x}^{j}\right) \right). \end{aligned}$$

$$(A.7)$$

Equation (A.2) can be rewritten as,

$$f(\mathbf{x}^{j+1}) - f(\mathbf{x}^{j}) \geq \frac{c}{N} \sum_{i=1}^{N} \frac{1}{h_{i}^{3}} k' \left(\left\| \frac{\mathbf{d}(\mathbf{l}_{i}, \mathbf{x}^{j})}{h} \right\|^{2} \right) \left[\left\| \mathbf{d}(\mathbf{l}_{i}, \mathbf{x}^{j+1}) \right\|^{2} - \left\| \mathbf{d}(\mathbf{l}_{i}, \mathbf{x}^{j}) \right\|^{2} \right]$$

by Inequality (A.4) (A.8)

$$\geq \frac{2c}{N} \sum_{i=1}^{N} \frac{1}{h_i^3} k' \left(\left\| \frac{\mathbf{d}(\mathbf{l}_i, \mathbf{x}^j)}{h} \right\|^2 \right) \mathbf{d}(\mathbf{l}_i, \mathbf{x}^j)^{\mathsf{T}} \left(\nabla_{\mathbf{x}} \mathbf{d}(\mathbf{l}_i, \mathbf{x}^j) \right) \left(\mathbf{x}^{j+1} - \mathbf{x}^j \right)$$

by Inequality (A.7) (A.9)

$$= \frac{2c}{N} \sum_{i=1}^{N} w_i^j \left(\widetilde{\mathbf{x}}_i^j - \mathbf{x}^j \right)^{\mathsf{T}} \left(\mathbf{x}^{j+1} - \mathbf{x}^j \right) \quad \text{by Equation (5)}$$
(A.10)

$$= \frac{2c}{N} \left[\left(\sum_{i=1}^{N} w_i^j (\widetilde{\mathbf{x}}_i^j)^\mathsf{T} \right) \mathbf{x}^{j+1} - \left(\sum_{i=1}^{N} w_i^j (\widetilde{\mathbf{x}}_i^j)^\mathsf{T} \right) \mathbf{x}^j \right]$$
(A.11)

$$-\sum_{i=1}^{N} w_i^j (\mathbf{x}^j)^{\mathsf{T}} \mathbf{x}^{j+1} + \sum_{i=1}^{N} w_i^j (\mathbf{x}^j)^{\mathsf{T}} \mathbf{x}^j \right] \qquad \text{by expansion} \qquad (A.12)$$

$$= \frac{2c}{N} \sum_{i=1}^{N} w_i^j \left(\|\mathbf{x}^{j+1}\|^2 - 2\left(\mathbf{x}^{j+1}\right)^\mathsf{T} \mathbf{x}^j + \|\mathbf{x}^j\|^2 \right)$$

because $\sum_{i=1}^{N} w_i^j \mathbf{x}^{j+1} = \sum_{i=1}^{N} w_i^j \widetilde{\mathbf{x}}_i^j$ from Equation (6) (A.13)

$$= \frac{2c}{N} \|\mathbf{x}^{j+1} - \mathbf{x}^{j}\|^{2} \sum_{i=1}^{N} w_{i}^{j}.$$
 (A.14)

Since the profile, k(x), is monotonically decreasing, k'(x) < 0. This leads the weight w_i^j to be strictly positive. As a result, the right hand side of Inequality (A.14) is strictly positive if $\mathbf{x}^j \neq \mathbf{x}^{j+1}$. Thus, $f(\mathbf{x}^{j+1}) - f(\mathbf{x}^j) > 0$.

References

- [1] H. Hindi. A tutorial on convex optimization. In American Control Conference, 2004.
- [2] S. Boyd and L. Vandenberghe. Convex Optimization. Cambridge University Press, 2004.