Algorithmic Regularization in Learning Deep Homogeneous Models: Layers are Automatically Balanced˚

Simon S. Du[†] Wei Hu^{\ddagger} Jason D. Lee[§]

Abstract

We study the implicit regularization imposed by gradient descent for learning multi-layer homogeneous functions including feed-forward fully connected and convolutional deep neural networks with linear, ReLU or Leaky ReLU activation. We rigorously prove that gradient flow (i.e. gradient descent with infinitesimal step size) effectively enforces the differences between squared norms across different layers to remain *invariant* without any explicit regularization. This result implies that if the weights are initially small, gradient flow automatically balances the magnitudes of all layers. Using a discretization argument, we analyze gradient descent with positive step size for the non-convex low-rank asymmetric matrix factorization problem without any regularization. Inspired by our findings for gradient flow, we prove that gradient descent with step sizes $\eta_t = O\left(t^{-\left(\frac{1}{2} + \delta\right)}\right)$ $(0 < \delta \leq \frac{1}{2})$ automatically balances two low-rank factors and converges to a bounded global optimum. Furthermore, for rank-1 asymmetric matrix factorization we give a finer analysis showing gradient descent with constant step size converges to the global minimum at a globally linear rate. We believe that the idea of examining the invariance imposed by first order algorithms in learning homogeneous models could serve as a fundamental building block for studying optimization for learning deep models.

1 Introduction

Modern machine learning models often consist of multiple layers. For example, consider a feedforward deep neural network that defines a prediction function

$$
\boldsymbol{x} \mapsto f(\boldsymbol{x}; \boldsymbol{W}^{(1)}, \dots, \boldsymbol{W}^{(N)}) = \boldsymbol{W}^{(N)} \phi(\boldsymbol{W}^{(N-1)} \cdots \boldsymbol{W}^{(2)} \phi(\boldsymbol{W}^{(1)} \boldsymbol{x}) \cdots),
$$

where $W^{(1)}, \ldots, W^{(N)}$ are weight matrices in N layers, and $\phi(\cdot)$ is a point-wise *homogeneous* activation function such as Rectified Linear Unit (ReLU) $\phi(x) = \max\{x, 0\}$. A simple observation is that this model is *homogeneous*: if we multiply a layer by a positive scalar c and divide another layer by c, the prediction function remains the same, e.g. $f(x; cW^{(1)}, \dots, \frac{1}{c}W^{(N)}) =$ $f(\boldsymbol{x};\boldsymbol{W}^{(1)},\dots,\boldsymbol{W}^{(N)}).$

A direct consequence of homogeneity is that a solution can produce small function value while being unbounded, because one can always multiply one layer by a huge number and divide another

32nd Conference on Neural Information Processing Systems (NeurIPS 2018), Montreal, Canada. ´

[˚]The full version of this paper is available at <https://arxiv.org/abs/1806.00900>.

[†]Machine Learning Department, School of Computer Science, Carnegie Mellon University. Email: ssdu@cs.cmu.edu

[;]Computer Science Department, Princeton University. Email: huwei@cs.princeton.edu

[§]Department of Data Sciences and Operations, Marshall School of Business, University of Southern California. Email: jasonlee@marshall.usc.edu

layer by that number. Theoretically, this possible unbalancedness poses significant difficulty in analyzing first order optimization methods like gradient descent/stochastic gradient descent (GD/SGD), because when parameters are not a priori constrained to a compact set via either coerciveness^{[5](#page-1-0)} of the loss or an explicit constraint, GD and SGD are not even guaranteed to converge [\[Lee et al.,](#page-10-0) [2016,](#page-10-0) Proposition 4.11]. In the context of deep learning, [Shamir](#page-10-1) [\[2018\]](#page-10-1) determined that the primary barrier to providing algorithmic results is in that the sequence of parameter iterates is possibly unbounded.

Now we take a closer look at asymmetric matrix factorization, which is a simple two-layer homogeneous model. Consider the following formulation for factorizing a low-rank matrix:

$$
\min_{\bm{U}\in\mathbb{R}^{d_1\times r}, \bm{V}\in\mathbb{R}^{d_2\times r}} f\left(\bm{U},\bm{V}\right) = \frac{1}{2} \left\| \bm{U}\bm{V}^{\top} - \bm{M}^* \right\|_F^2, \tag{1}
$$

where $M^* \in \mathbb{R}^{d_1 \times d_2}$ is a matrix we want to factorize. We observe that due to the homogeneity of f , it is not smooth^{[6](#page-1-1)} even in the neighborhood of a globally optimum point. To see this, we compute the gradient of f :

$$
\frac{\partial f(\boldsymbol{U}, \boldsymbol{V})}{\partial \boldsymbol{U}} = (\boldsymbol{U}\boldsymbol{V}^\top - \boldsymbol{M}^*) \boldsymbol{V}, \qquad \frac{\partial f(\boldsymbol{U}, \boldsymbol{V})}{\partial \boldsymbol{V}} = (\boldsymbol{U}\boldsymbol{V}^\top - \boldsymbol{M}^*)^\top \boldsymbol{U}.
$$
 (2)

Notice that the gradient of f is not homogeneous anymore. Further, consider a globally optimal solution (U, V) such that $||U||_F$ is of order ϵ and $||V||_F$ is of order $1/\epsilon$ (ϵ being very small). A small perturbation on U can lead to dramatic change to the gradient of U . This phenomenon can happen for all homogeneous functions when the layers are unbalanced. The lack of nice geometric properties of homogeneous functions due to unbalancedness makes first-order optimization methods difficult to analyze.

A common theoretical workaround is to artificially modify the natural objective function as in [\(1\)](#page-1-2) in order to prove convergence. In [\[Tu et al.,](#page-10-2) [2015,](#page-10-2) [Ge et al.,](#page-9-0) [2017a\]](#page-9-0), a regularization term for balancing the two layers is added to (1) :

$$
\min_{\bm{U}\in\mathbb{R}^{d_1\times r},\bm{V}\in\mathbb{R}^{d_2\times r}}\frac{1}{2}\left\|\bm{U}\bm{V}^{\top}-\bm{M}\right\|_{F}^{2}+\frac{1}{8}\left\|\bm{U}^{\top}\bm{U}-\bm{V}^{\top}\bm{V}\right\|_{F}^{2}.
$$
\n(3)

For problem [\(3\)](#page-1-3), the regularizer removes the homogeneity issue and the optimal solution becomes unique (up to rotation). [Ge et al.](#page-9-0) $[2017a]$ showed that the modified objective (3) satisfies (i) every local minimum is a global minimum, (ii) all saddle points are strict^{[7](#page-1-4)}, and (iii) the objective is smooth. These imply that (noisy) GD finds a global minimum [\[Ge et al.,](#page-9-1) [2015,](#page-9-1) [Lee et al.,](#page-10-0) [2016,](#page-10-0) [Panageas](#page-10-3) [and Piliouras,](#page-10-3) [2016\]](#page-10-3).

On the other hand, empirically, removing the homogeneity is not necessary. We use GD with random initialization to solve the optimization problem [\(1\)](#page-1-2). Figure [1a](#page-2-0) shows that even *without* regularization term like in the modified objective [\(3\)](#page-1-3) GD with random initialization converges to a global minimum and the convergence rate is also competitive. A more interesting phenomenon is shown in Figure [1b](#page-2-0) in which we track the Frobenius norms of U and V in all iterations. The plot shows that the ratio between norms remains a constant in all iterations. Thus the unbalancedness does not occur at all! In many practical applications, many models also admit the homogeneous property (like deep neural networks) and first order methods often converge to a balanced solution. A natural question arises:

Why does GD balance multiple layers and converge in learning homogeneous functions?

In this paper, we take an important step towards answering this question. Our key finding is that the gradient descent algorithm provides an implicit regularization on the target homogeneous function. First, we show that on the gradient flow (gradient descent with infinitesimal step size) trajectory induced by any differentiable loss function, for a large class of homogeneous models, including fully connected and convolutional neural networks with linear, ReLU and Leaky ReLU activations, the differences between squared norms across layers remain invariant. Thus, as long as at the beginning the differences are small, they remain small at all time. Note that small differences arise in commonly used initialization schemes such as $\frac{1}{\sqrt{2}}$ \overline{d} Gaussian initialization or

⁵A function f is coercive if $||x|| \to \infty$ implies $f(x) \to \infty$.

⁶A function is said to be smooth if its gradient is β -Lipschitz continuous for some finite $\beta > 0$.

 7 A saddle point of a function f is strict if the Hessian at that point has a negative eigenvalue.

(a) Comparison of convergence rates of GD for objective functions (1) and (3) .

(b) Comparison of quantity $||\boldsymbol{U}||_F^2 / ||\boldsymbol{V}||_F^2$ when running GD for objective functions (1) and (3) .

Figure 1: Experiments on the matrix factorization problem with objective functions [\(1\)](#page-1-2) and [\(3\)](#page-1-3). Red lines correspond to running GD on the objective function [\(1\)](#page-1-2), and blue lines correspond to running GD on the objective function (3) .

Xavier/Kaiming initialization schemes [\[Glorot and Bengio,](#page-9-2) [2010,](#page-9-2) [He et al.,](#page-10-4) [2016\]](#page-10-4). Our result thus explains why using ReLU activation is a better choice than sigmoid from the optimization point view. For linear activation, we prove an even stronger invariance for gradient flow: we show that $W^{(h)} (W^{(h)})^{\top} - (W^{(h+1)})^{\top} W^{(h+1)}$ stays invariant over time, where $W^{(h)}$ and $W^{(h+1)}$ are weight matrices in consecutive layers with linear activation in between.

Next, we go beyond gradient flow and consider gradient descent with positive step size. We focus on the asymmetric matrix factorization problem [\(1\)](#page-1-2). Our invariance result for linear activation indicates that $U^\top U - V^\top V$ stays unchanged for gradient flow. For gradient descent, $U^\top U - V^\top V$ can change over iterations. Nevertheless we show that if the step size decreases like $\eta_t = O\left(t^{-\left(\frac{1}{2}+\delta\right)}\right)$ $(0 < \delta \leq \frac{1}{2})$, $\mathbf{U}^\top \mathbf{U} - \mathbf{V}^\top \mathbf{V}$ will remain small in all iterations. In the set where $\mathbf{U}^\top \mathbf{U} - \mathbf{V}^\top \mathbf{V}$ is small, the loss is coercive, and gradient descent thus ensures that all the iterates are bounded. Using these properties, we then show that gradient descent converges to a globally optimal solution. Furthermore, for rank-1 asymmetric matrix factorization, we give a finer analysis and show that randomly initialized gradient descent with *constant* step size converges to the global minimum at a globally linear rate.

Related work. The homogeneity issue has been previously discussed by [Neyshabur et al.](#page-10-5) [\[2015a,](#page-10-5)[b\]](#page-10-6). The authors proposed a variant of stochastic gradient descent that regularizes paths in a neural network, which is related to the max-norm. The algorithm outperforms gradient descent and AdaGrad on several classification tasks.

A line of research focused on analyzing gradient descent dynamics for (convolutional) neural networks with one or two unknown layers [\[Tian,](#page-10-7) [2017,](#page-10-7) [Brutzkus and Globerson,](#page-9-3) [2017,](#page-9-3) [Du et al.,](#page-9-4) [2017a](#page-9-4)[,b,](#page-9-5) [Zhong et al.,](#page-11-0) [2017,](#page-11-0) [Li and Yuan,](#page-10-8) [2017,](#page-10-8) [Ma et al.,](#page-10-9) [2017,](#page-10-9) [Brutzkus et al.,](#page-9-6) [2017\]](#page-9-6). For one unknown layer, there is no homogeneity issue. While for two unknown layers, existing work either re-quires learning two layers separately [\[Zhong et al.,](#page-11-0) [2017,](#page-11-0) [Ge et al.,](#page-9-7) [2017b\]](#page-9-7) or uses re-parametrization like weight normalization to remove the homogeneity issue [\[Du et al.,](#page-9-5) [2017b\]](#page-9-5). To our knowledge, there is no rigorous analysis for optimizing multi-layer homogeneous functions.

For a general (non-convex) optimization problem, it is known that if the objective function satisfies (i) gradient changes smoothly if the parameters are perturbed, (ii) all saddle points and local maxima are strict (i.e., there exists a direction with negative curvature), and (iii) all local minima are global (no spurious local minimum), then gradient descent [\[Lee et al.,](#page-10-0) [2016,](#page-10-0) [Panageas and Piliouras,](#page-10-3) [2016\]](#page-10-3) converges to a global minimum. There have been many studies on the optimization landscapes of neural networks [\[Kawaguchi,](#page-10-10) [2016,](#page-10-10) [Choromanska et al.,](#page-9-8) [2015,](#page-9-8) [Du and Lee,](#page-9-9) [2018,](#page-9-9) [Hardt and](#page-10-11) [Ma,](#page-10-11) [2016,](#page-10-11) [Bartlett et al.,](#page-9-10) [2018,](#page-9-10) [Haeffele and Vidal,](#page-10-12) [2015,](#page-10-12) [Freeman and Bruna,](#page-9-11) [2016,](#page-9-11) [Vidal et al.,](#page-10-13) [2017,](#page-10-13) [Safran and Shamir,](#page-10-14) [2016,](#page-10-14) [Zhou and Feng,](#page-11-1) [2017,](#page-11-1) [Nguyen and Hein,](#page-10-15) [2017a](#page-10-15)[,b,](#page-10-16) [Zhou and Feng,](#page-11-1) [2017,](#page-11-1) [Safran and Shamir,](#page-10-17) [2017\]](#page-10-17), showing that the objective functions have properties (ii) and (iii).

Nevertheless, the objective function is in general not smooth as we discussed before. Our paper complements these results by showing that the magnitudes of all layers are balanced and in many cases, this implies smoothness.

Paper organization. The rest of the paper is organized as follows. In Section [2,](#page-3-0) we present our main theoretical result on the implicit regularization property of gradient flow for optimizing neural networks. In Section [3,](#page-6-0) we analyze the dynamics of randomly initialized gradient descent for asymmetric matrix factorization problem with unregularized objective function [\(1\)](#page-1-2). In Section [4,](#page-7-0) we empirically verify the theoretical result in Section [2.](#page-3-0) We conclude and list future directions in Section [5.](#page-8-0) Some technical proofs are deferred to the appendix.

Notation. We use bold-faced letters for vectors and matrices. For a vector x, denote by $x[i]$ its *i*-th coordinate. For a matrix A, we use $A[i, j]$ to denote its (i, j) -th entry, and use $A[i, :]$ and $A[:, j]$ to denote its *i*-th row and *j*-th column, respectively (both as column vectors). We use $\|\cdot\|_2$ or $\|\cdot\|$ to denote the Euclidean norm of a vector, and use $\|\cdot\|_F$ to denote the Frobenius norm of a matrix. We use $\langle \cdot, \cdot \rangle$ to denote the standard Euclidean inner product between two vectors or two matrices. Let $[n] = \{1, 2, \ldots, n\}.$

2 The Auto-Balancing Properties in Deep Neural Networks

In this section we study the implicit regularization imposed by gradient descent with infinitesimal step size (gradient flow) in training deep neural networks. In Section [2.1](#page-3-1) we consider fully connected neural networks, and our main result (Theorem [2.1\)](#page-4-0) shows that gradient flow automatically balances the incoming and outgoing weights at every neuron. This directly implies that the weights between different layers are balanced (Corollary [2.1\)](#page-4-1). For linear activation, we derive a stronger auto-balancing property (Theorem [2.2\)](#page-4-2). In Section [2.2](#page-4-3) we generalize our result from fully connected neural networks to convolutional neural networks. In Section [2.3](#page-5-0) we present the proof of Theorem [2.1.](#page-4-0) The proofs of other theorems in this section follow similar ideas and are deferred to Appendix [A.](#page-12-0)

2.1 Fully Connected Neural Networks

We first formally define a fully connected feed-forward neural network with $N (N \ge 2)$ layers. Let $W^{(h)} \in \mathbb{R}^{n_h \times n_{h-1}}$ be the weight matrix in the h-th layer, and define $w = (W^{(h)})_{h=1}^N$ as a shorthand of the collection of all the weights. Then the function $f_w : \mathbb{R}^d \to \mathbb{R}^p$ ($d = n_0, p = n_N$) computed by this network can be defined recursively: $f_{w}^{(1)}(x) = W^{(1)}x$, $f_{w}^{(h)}(x) = W^{(h)}\phi_{h-1}(f_{w}^{(h-1)}(x))$ $(h = 2, \ldots, N)$, and $f_{\bm{w}}(\bm{x}) = f_{\bm{w}}^{(N)}(\bm{x})$, where each ϕ_h is an activation function that acts coordinate-wise on vectors.^{[8](#page-3-2)} We assume that each ϕ_h ($h \in [N-1]$) is *homogeneous*, namely, $\phi_h(x) = \phi'_h(x) \cdot x$ for all x and all elements of the sub-differential $\phi'_h(\cdot)$ when ϕ_h is non-differentiable at x. This property is satisfied by functions like ReLU $\phi(x) = \max\{x, 0\}$, Leaky ReLU $\phi(x) = \max\{x, \alpha x\}$ $(0 < \alpha < 1)$, and linear function $\phi(x) = x$.

Let $\ell : \mathbb{R}^p \times \mathbb{R}^p \to \mathbb{R}_{\geqslant 0}$ be a differentiable loss function. Given a training dataset $\{(x_i, y_i)\}_{i=1}^m$ $\mathbb{R}^d \times \mathbb{R}^p$, the training loss as a function of the network parameters \bm{w} is defined as

$$
L(\boldsymbol{w}) = \frac{1}{m} \sum_{i=1}^{m} \ell \left(f_{\boldsymbol{w}}(\boldsymbol{x}_i), \boldsymbol{y}_i \right). \tag{4}
$$

We consider gradient descent with infinitesimal step size (also known as gradient flow) applied on $L(\boldsymbol{w})$, which is captured by the differential inclusion:

$$
\frac{\mathrm{d} \boldsymbol{W}^{(h)}}{\mathrm{d} t} \in -\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h)}}, \qquad h = 1, \dots, N,
$$
\n(5)

where t is a continuous time index, and $\frac{\partial L(w)}{\partial \mathbf{W}^{(h)}}$ is the Clarke sub-differential [\[Clarke et al.,](#page-9-12) [2008\]](#page-9-12). If curves $\mathbf{W}^{(h)} = \mathbf{W}^{(h)}(t)$ ($h \in [N]$) evolve with time according to [\(5\)](#page-3-3) they are said to be a solution of the gradient flow differential inclusion.

⁸We omit the trainable bias weights in the network for simplicity, but our results can be directly generalized to allow bias weights.

Our main result in this section is the following invariance imposed by gradient flow.

Theorem 2.1 (Balanced incoming and outgoing weights at every neuron). *For any* $h \in [N-1]$ *and* $i \in [n_h]$, we have

$$
\frac{\mathrm{d}}{\mathrm{d}t}\left(\|\mathbf{W}^{(h)}[i,:]\|^2 - \|\mathbf{W}^{(h+1)}[:,i]\|^2\right) = 0.
$$
\n(6)

Note that $\bm{W}^{(h)}[i,:]$ is a vector consisting of network weights coming into the i -th neuron in the h -th hidden layer, and $W^{(h+1)}[:, i]$ is the vector of weights going out from the same neuron. Therefore, Theorem [2.1](#page-4-0) shows that gradient flow exactly preserves the difference between the squared ℓ_2 -norms of incoming weights and outgoing weights at any neuron.

Taking sum of [\(6\)](#page-4-4) over $i \in [n_h]$, we obtain the following corollary which says gradient flow preserves the difference between the squares of Frobenius norms of weight matrices.

Corollary 2.1 (Balanced weights across layers). *For any* $h \in [N - 1]$, we have

$$
\frac{\mathrm{d}}{\mathrm{d}t}\left(\|\mathbf{W}^{(h)}\|_F^2 - \|\mathbf{W}^{(h+1)}\|_F^2\right) = 0.
$$

Corollary [2.1](#page-4-1) explains why in practice, trained multi-layer models usually have similar magnitudes on all the layers: if we use a small initialization, $\|\mathbf{W}^{(h)}\|_F^2 - \|\mathbf{W}^{(h+1)}\|_F^2$ is very small at the beginning, and Corollary [2.1](#page-4-1) implies this difference remains small at all time. This finding also partially explains why gradient descent converges. Although the objective function like [\(4\)](#page-3-4) may not be smooth over the entire parameter space, given that $\|{\bm{W}}^{(h)}\|_F^2 - \|{\bm{W}}^{(h+1)}\|_F^2$ is small for all h , the objective function may have smoothness. Under this condition, standard theory shows that gradient descent converges. We believe this finding serves as a key building block for understanding first order methods for training deep neural networks.

For linear activation, we have the following stronger invariance than Theorem [2.1:](#page-4-0)

Theorem 2.2 (Stronger balancedness property for linear activation). *If for some* $h \in [N - 1]$ *we have* $\phi_h(x) = x$ *, then*

$$
\frac{\mathrm{d}}{\mathrm{d} t}\left(\boldsymbol{W}^{(h)}(\boldsymbol{W}^{(h)})^\top - (\boldsymbol{W}^{(h+1)})^\top \boldsymbol{W}^{(h+1)}\right) = \boldsymbol{0}.
$$

This result was known for linear networks [\[Arora et al.,](#page-9-13) [2018\]](#page-9-13), but the proof there relies on the entire network being linear while Theorem [2.2](#page-4-2) only needs two consecutive layers to have no nonlinear activations in between.

While Theorem [2.1](#page-4-0) shows the invariance in a node-wise manner, Theorem [2.2](#page-4-2) shows for linear activation, we can derive a layer-wise invariance. Inspired by this strong invariance, in Section [3](#page-6-0) we prove gradient descent with positive step sizes preserves this invariance approximately for matrix factorization.

2.2 Convolutional Neural Networks

Now we show that the conservation property in Corollary [2.1](#page-4-1) can be generalized to convolutional neural networks. In fact, we can allow *arbitrary sparsity pattern and weight sharing structure* within a layer; convolutional layers are a special case.

Neural networks with sparse connections and shared weights. We use the same notation as in Section [2.1,](#page-3-1) with the difference that some weights in a layer can be *missing* or *shared*. Formally, the weight matrix $W^{(h)} \in \mathbb{R}^{n_h \times n_{h-1}}$ in layer h ($h \in [N]$) can be described by a vector $v^{(h)} \in \mathbb{R}^{d_h}$ and a function $g_h: [n_h] \times [n_{h-1}] \to [d_h] \cup \{0\}$. Here $v^{(h)}$ consists of the actual *free parameters* in this layer and d_h is the number of free parameters (e.g. if there are k convolutional filters in layer h each with size r, we have $d_h = r \cdot k$). The map g_h represents the sparsity and weight sharing pattern:

$$
\mathbf{W}^{(h)}[i,j] = \begin{cases} 0, & g_h(i,j) = 0, \\ \mathbf{v}^{(h)}[k], & g_h(i,j) = k > 0. \end{cases}
$$

Denote by $v =$ $\bm{v}^{(h)}\big)_{h_1}^N$ $h_{h=1}$ the collection of all the parameters in this network, and we consider gradient flow to learn the parameters:

$$
\frac{\mathrm{d}\boldsymbol{v}^{(h)}}{\mathrm{d}t}\in -\frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{v}^{(h)}}, \qquad h=1,\ldots,N.
$$

The following theorem generalizes Corollary [2.1](#page-4-1) to neural networks with sparse connections and shared weights:

Theorem 2.3. *For any* $h \in [N - 1]$ *, we have*

$$
\frac{\mathrm{d}}{\mathrm{d}t}\left(\|\boldsymbol{v}^{(h)}\|^2-\|\boldsymbol{v}^{(h+1)}\|^2\right)=0.
$$

Therefore, for a neural network with arbitrary sparsity pattern and weight sharing structure, gradient flow still balances the magnitudes of all layers.

2.3 Proof of Theorem [2.1](#page-4-0)

The proofs of all theorems in this section are similar. They are based on the use of the chain rule (i.e. back-propagation) and the property of homogeneous activations. Below we provide the proof of Theorem [2.1](#page-4-0) and defer the proofs of other theorems to Appendix [A.](#page-12-0)

Proof of Theorem [2.1.](#page-4-0) First we note that we can without loss of generality assume L is the loss associated with one data sample $(x, y) \in \mathbb{R}^d \times \mathbb{R}^p$, i.e., $L(w) = \ell(f_w(x), y)$. In fact, for $L(\boldsymbol{w}) = \frac{1}{m}$ vith
マ $_{k=1}^{m} L_k(\boldsymbol{w})$ where $L_k(\boldsymbol{w}) = \ell(f_{\boldsymbol{w}}(\boldsymbol{x}_k), \boldsymbol{y}_k)$, for any single weight $\boldsymbol{W}^{(h)}[i, j]$ in the network we can compute $\frac{\mathrm{d}}{\mathrm{d}t}(\boldsymbol{W}^{(h)}[i,j])^2 = 2\boldsymbol{W}^{(h)}[i,j] \cdot \frac{\mathrm{d}\boldsymbol{W}^{(h)}[i,j]}{\mathrm{d}t} = -2\boldsymbol{W}^{(h)}[i,j] \cdot \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h)}[i,j]} =$ $-2{\boldsymbol W}^{(h)}[i,j] \cdot \frac{1}{m}$ $\sum_{n=1}^{\infty}$ $\frac{\partial L_k(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h)}[i]_n}$ $\frac{\partial L_k(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h)}[i,j]}$, using the sharp chain rule of differential inclusions for tame functions [\[Drusvyatskiy et al.,](#page-9-14) [2015,](#page-9-14) [Davis et al.,](#page-9-15) [2018\]](#page-9-15). Thus, if we can prove the theorem for every individual loss L_k , we can prove the theorem for L by taking average over $k \in [m]$.

Therefore in the rest of proof we assume $L(\bm{w}) = \ell(f_{\bm{w}}(\bm{x}), \bm{y})$. For convenience, we denote $\bm{x}^{(h)}$ = $f_{\bm{w}}^{(h)}(\bm{x})$ ($h \in [N]$), which is the input to the h-th hidden layer of neurons for $h \in [N-1]$ and is the output of the network for $h = N$. We also denote $\boldsymbol{x}^{(0)} = \boldsymbol{x}$ and $\phi_0(x) = x$ ($\forall x$).

Now we prove [\(6\)](#page-4-4). Since $W^{(h+1)}[k, i]$ ($k \in [n_{h+1}]$) can only affect $L(w)$ through $x^{(h+1)}[k]$, we have for $k \in [n_{h+1}],$

$$
\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h+1)}[k,i]} = \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}[k]} \cdot \frac{\partial \boldsymbol{x}^{(h+1)}[k]}{\partial \boldsymbol{W}^{(h+1)}[k,i]} = \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}[k]} \cdot \phi_h(\boldsymbol{x}^{(h)}[i]),
$$

which can be rewritten as

$$
\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h+1)}[:,i]} = \phi_h(\boldsymbol{x}^{(h)}[i]) \cdot \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}}.
$$

It follows that

$$
\frac{\mathrm{d}}{\mathrm{d}t} \|\mathbf{W}^{(h+1)}[:,i]\|^2 = 2 \left\langle \mathbf{W}^{(h+1)}[:,i],\frac{\mathrm{d}}{\mathrm{d}t} \mathbf{W}^{(h+1)}[:,i] \right\rangle = -2 \left\langle \mathbf{W}^{(h+1)}[:,i],\frac{\partial L(\mathbf{w})}{\partial \mathbf{W}^{(h+1)}[:,i]} \right\rangle
$$
\n
$$
= -2\phi_h(\mathbf{x}^{(h)}[i]) \cdot \left\langle \mathbf{W}^{(h+1)}[:,i],\frac{\partial L(\mathbf{w})}{\partial \mathbf{x}^{(h+1)}} \right\rangle. \tag{7}
$$

On the other hand, $W^{(h)}[i, :]$ only affects $L(w)$ through $x^{(h)}[i]$. Using the chain rule, we get F

$$
\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h)}[i, :]} = \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h)}[i]} \cdot \phi_{h-1}(\boldsymbol{x}^{(h-1)}) = \left\langle \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}}, \boldsymbol{W}^{(h+1)}[:, i] \right\rangle \cdot \phi'_{h}(\boldsymbol{x}^{(h)}[i]) \cdot \phi_{h-1}(\boldsymbol{x}^{(h-1)}),
$$

where ϕ' is interpreted as a set-valued mapping whenever it is applied at a non-differentiable point.^{[9](#page-5-1)}

⁹More precisely, the equalities should be an inclusion whenever there is a sub-differential, but as we see in the next display the ambiguity in the choice of sub-differential does not affect later calculations.

It follows that 10

$$
\frac{\mathrm{d}}{\mathrm{d}t} \|\mathbf{W}^{(h)}[i, :] \|^2 = 2 \left\langle \mathbf{W}^{(h)}[i, :] , \frac{\mathrm{d}}{\mathrm{d}t} \mathbf{W}^{(h)}[i, :] \right\rangle = -2 \left\langle \mathbf{W}^{(h)}[i, :] , \frac{\partial L(\mathbf{w})}{\partial \mathbf{W}^{(h)}[i, :]} \right\rangle
$$
\n
$$
= -2 \left\langle \frac{\partial L(\mathbf{w})}{\partial \mathbf{x}^{(h+1)}}, \mathbf{W}^{(h+1)}[:, i] \right\rangle \cdot \phi'_h(\mathbf{x}^{(h)}[i]) \cdot \left\langle \mathbf{W}^{(h)}[i, :] , \phi_{h-1}(\mathbf{x}^{(h-1)}) \right\rangle
$$
\n
$$
= -2 \left\langle \frac{\partial L(\mathbf{w})}{\partial \mathbf{x}^{(h+1)}}, \mathbf{W}^{(h+1)}[:, i] \right\rangle \cdot \phi'_h(\mathbf{x}^{(h)}[i]) \cdot \mathbf{x}^{(h)}[i] = -2 \left\langle \frac{\partial L(\mathbf{w})}{\partial \mathbf{x}^{(h+1)}}, \mathbf{W}^{(h+1)}[:, i] \right\rangle \cdot \phi_h(\mathbf{x}^{(h)}[i]).
$$

Comparing the above expression to [\(7\)](#page-5-2), we finish the proof.

3 Gradient Descent Converges to Global Minimum for Asymmetric Matrix Factorization

In this section we constrain ourselves to the asymmetric matrix factorization problem and analyze the gradient descent algorithm with random initialization. Our analysis is inspired by the auto-balancing properties presented in Section [2.](#page-3-0) We extend these properties from gradient flow to gradient descent with positive step size.

Formally, we study the following non-convex optimization problem:

$$
\min_{\bm{U} \in \mathbb{R}^{d_1 \times r}, \bm{V} \in \mathbb{R}^{d_2 \times r}} f(\bm{U}, \bm{V}) = \frac{1}{2} \left\| \bm{U} \bm{V}^\top - \bm{M}^* \right\|_F^2, \tag{8}
$$

 \Box

where $M^* \in \mathbb{R}^{d_1 \times d_2}$ has rank r. Note that we do not have any explicit regularization in [\(8\)](#page-6-2). The gradient descent dynamics for [\(8\)](#page-6-2) have the following form:

$$
U_{t+1} = U_t - \eta_t (U_t V_t^\top - M^*) V_t, \qquad V_{t+1} = V_t - \eta_t (U_t V_t^\top - M^*)^\top U_t.
$$
 (9)

3.1 The General Rank-r Case

First we consider the general case of $r \geq 1$. Our main theorem below says that if we use a random small initialization (U_0, V_0) , and set step sizes η_t to be appropriately small, then gradient descent [\(9\)](#page-6-3) will converge to a solution close to the global minimum of [\(8\)](#page-6-2). To our knowledge, this is the first result showing that gradient descent with random initialization directly solves the un-regularized asymmetric matrix factorization problem [\(8\)](#page-6-2).

Theorem 3.1. Let $0 < \epsilon < \|M^*\|_F$. Suppose we initialize the entries in U_0 and V_0 i.i.d. from $\mathcal{N}(0,\frac{\epsilon}{\text{poly}(d)})$ ($d=\max\{d_1,d_2\}$), and run [\(9\)](#page-6-3) with step sizes $\eta_t=\frac{\sqrt{\epsilon/r}}{100(t+1)\|{\bm{M}}^2}$ $\frac{\sqrt{\epsilon/r}}{100(t+1)\|\mathbf{M}^*\|_F^{3/2}}$ $(t = 0, 1, \ldots).^{11}$ $(t = 0, 1, \ldots).^{11}$ $(t = 0, 1, \ldots).^{11}$ *Then with high probability over the initialization,* $\lim_{t\to\infty} (U_t, V_t) = (\tilde{U}, \tilde{V})$ *exists and satisfies* $\left\| \bar{U} \bar{V}^\top - M^* \right\|_F \leqslant \epsilon.$

Proof sketch of Theorem [3.1.](#page-6-5) First let's imagine that we are using infinitesimal step size in GD. Then according to Theorem [2.2](#page-4-2) (viewing problem [\(8\)](#page-6-2) as learning a two-layer linear network where the inputs are all the standard unit vectors in \mathbb{R}^{d_2}), we know that $U^{\top}U - V^{\top}V$ will stay invariant throughout the algorithm. Hence when U and V are initialized to be small, $U^\top U - V^\top V$ will stay small forever. Combined with the fact that the objective $f(U,V)$ is decreasing over time (which means UV^{\top} cannot be too far from M^*), we can show that U and V will always stay bounded.

Now we are using positive step sizes η_t , so we no longer have the invariance of $\bm{U}^\top \bm{U} - \bm{V}^\top \bm{V}$. Nevertheless, by a careful analysis of the updates, we can still prove that $U_t^\top U_t - V_t^\top V_t$ is small, the objective $f(U_t, V_t)$ decreases, and U_t and V_t stay bounded. Formally, we have the following lemma:

Lemma 3.1. *With high probability over the initialization* (U_0, V_0) *, for all t we have:*

¹⁰This holds for any choice of element of the sub-differential, since $\phi'(x)x = \phi(x)$ holds at $x = 0$ for any choice of sub-differential.

¹¹The dependency of η_t on t can be $\eta_t = \Theta$ $t^{-(1/2+\delta)}$ for any constant $\delta \in (0, 1/2]$.

- *(i) Balancedness:* $\begin{aligned} \big\| \boldsymbol{U}_t^\top \boldsymbol{U}_t - \boldsymbol{V}_t^\top \boldsymbol{V}_t \end{aligned}$ $\big\|_F \leqslant \epsilon;$
- *(ii)* Decreasing objective: $f(U_t, V_t) \leqslant f(U_{t-1}, V_{t-1}) \leqslant \cdots \leqslant f(U_0, V_0) \leqslant 2 \|M^*\|_F^2$;
- *(iii) Boundedness:* $||U_t||_F^2 \le 5\sqrt{r} ||M^*||_F$, $||V_t||_F^2 \le 5\sqrt{r} ||M^*||_F$.

Now that we know the GD algorithm automatically constrains (U_t, V_t) in a bounded region, we can use the smoothness of f in this region and a standard analysis of GD to show that (U_t, V_t) converges to a stationary point (\bar{U}, \bar{V}) of f (Lemma [B.2\)](#page-16-0). Furthermore, using the results of [\[Lee](#page-10-0) [et al.,](#page-10-0) [2016,](#page-10-0) [Panageas and Piliouras,](#page-10-3) [2016\]](#page-10-3) we know that (\bar{U}, \bar{V}) is almost surely not a strict saddle point. Then the following lemma implies that (\bar{U}, \bar{V}) has to be close to a global optimum since we know $\|\vec{U}^\top \vec{U} - \vec{V}^\top \vec{V}\|_F \le \epsilon$ from Lemma [3.1](#page-6-6) (i). This would complete the proof of Theorem [3.1.](#page-6-5)

Lemma 3.2. Suppose (U, V) is a stationary point of f such that $\left\|U^{\top}U - V^{\top}V\right\|_F \leq \epsilon$. Then *either* **a 5.2.** Suppose (U, V) is a stationary point of f such that $\left\| UV^\top - M^* \right\|_F \leq \epsilon$, or (U, V) is a strict saddle point of f.

The full proof of Theorem [3.1](#page-6-5) and the proofs of Lemmas [3.1](#page-6-6) and [3.2](#page-7-1) are given in Appendix [B.](#page-13-0)

3.2 The Rank-1 Case

We have shown in Theorem [3.1](#page-6-5) that GD with small and diminishing step sizes converges to a global minimum for matrix factorization. Empirically, it is observed that a constant step size $\eta_t \equiv \eta$ is enough for GD to converge quickly to global minimum. Therefore, some natural questions are how to prove convergence of GD with a constant step size, how fast it converges, and how the discretization affects the invariance we derived in Section [2.](#page-3-0)

While these questions remain challenging for the general rank- r matrix factorization, we resolve them for the case of $r = 1$. Our main finding is that with constant step size, the norms of two layers are always within a constant factor of each other (although we may no longer have the stronger balancedness property as in Lemma [3.1\)](#page-6-6), and we utilize this property to prove the *linear convergence* of GD to a global minimum.

When $r = 1$, the asymmetric matrix factorization problem and its GD dynamics become

$$
\min_{\bm{u}\in\mathbb{R}^{d_1},\bm{v}\in\mathbb{R}^{d_2}}\frac{1}{2}\left\Vert \bm{u}\bm{v}^\top - \bm{M}^*\right\Vert_F^2
$$

and

$$
\boldsymbol{u}_{t+1} = \boldsymbol{u}_t - \eta (\boldsymbol{u}_t \boldsymbol{v}_t^\top - \boldsymbol{M}^*) \boldsymbol{v}_t, \qquad \boldsymbol{v}_{t+1} = \boldsymbol{v}_t - \eta \left(\boldsymbol{v}_t \boldsymbol{u}_t^\top - \boldsymbol{M}^{*\top}\right) \boldsymbol{u}_t.
$$

Here we assume M^* has rank 1, i.e., it can be factorized as $M^* = \sigma_1 u^* v^{*\top}$ where u^* and v^* are unit vectors and $\sigma_1 > 0$.

Our main theoretical result is the following.

Theorem 3.2 (Approximate balancedness and linear convergence of GD for rank-1 matrix factor-**Theorem 3.2** (Approximate balancedness and linear convergence of GD for rank-1 matrix factorization). *Suppose* $u_0 \sim \mathcal{N}(0, \delta I)$, $v_0 \sim \mathcal{N}(0, \delta I)$ with $\delta = c_{init} \sqrt{\frac{\sigma_1}{d}} (d = \max\{d_1, d_2\})$ for some *sufficiently small constant* $c_{init} > 0$ *, and* $\eta = \frac{c_{step}}{\sigma_1}$ $\frac{step}{\sigma_1}$ for some sufficiently small constant $c_{step} > 0$. Then with constant probability over the initialization, for all t we have $c_0 \leq \frac{|u_t^\top u^*|}{|u^\top u^*|}$ $\left|\frac{u_t}{v_t^\top v^*}\right| \leq C_0$ for some *universal constants* $c_0, C_0 > 0$ *. Furthermore, for any* $0 < \epsilon < 1$ *, after* $t = O$ \boldsymbol{v} $\log \frac{d}{\epsilon}$ *iterations, we have* rsal constants $\| \bm{u}_t \bm{v}_t^{\top} - \bm{M^*}$ $c_0, C_0 > 0$
 $\Vert_F \leq \epsilon \sigma_1.$

Theorem [3.2](#page-7-2) shows for u_t and v_t , their strengths in the signal space, $|u_t^\top u^*|$ and $|v_t^\top v^*|$, are of the same order. This approximate balancedness helps us prove the linear convergence of GD. We refer readers to Appendix [C](#page-18-0) for the proof of Theorem [3.2.](#page-7-2)

4 Empirical Verification

We perform experiments to verify the auto-balancing properties of gradient descent in neural networks with ReLU activation. Our results below show that for GD with small step size and small

Figure 2: Balancedness of a 3-layer neural network.

initialization: (1) the difference between the squared Frobenius norms of any two layers remains small in all iterations, and (2) the ratio between the squared Frobenius norms of any two layers becomes close to 1. Notice that our theorems in Section [2](#page-3-0) hold for gradient flow (step size \rightarrow 0) but in practice we can only choose a (small) positive step size, so we cannot hope the difference between the squared Frobenius norms to remain exactly the same but can only hope to observe that the differences remain small.

We consider a 3-layer fully connected network of the form $f(x) = W_3\phi(W_2\phi(W_1x))$ where $x \in \mathbb{R}^{1,000}$ is the input, $W_1 \in \mathbb{R}^{100 \times 1,000}$, $W_2 \in \mathbb{R}^{100 \times 100}$, $W_3 \in \mathbb{R}^{10 \times 100}$, and $\phi(\cdot)$ is ReLU activation. We use 1,000 data points and the quadratic loss function, and run GD. We first test a balanced initialization: $W_1[i, j] \sim N(0, \frac{10^{-4}}{100})$, $W_2[i, j] \sim N(0, \frac{10^{-4}}{10})$ and $W_3[i, j] \sim N(0, 10^{-4})$, which ensures $||W_1||_F^2 \approx ||W_2||_F^2 \approx ||\dot{W}_3||_F^2$. After 10,000 iterations we have $||W_1||_F^2 = 42.90$,
 $||W_2||_F^2 = 43.76$ and $||W_2||_F^2 = 43.68$. Figure 23 shows that in all iterations $||W_1||_F^2 = ||W_2||_F^2$ $||W_2||_F^2 = 43.76$ and $||W_3||_F^2 = 43.68$. Figure [2a](#page-8-1) shows that in all iterations $||W_1||_F^2 - ||W_2||_F^2$ and $\|W_2\|_F^2 - \|W_3\|_F^2$ are bounded by 0.14 which is much smaller than the magnitude of each $||W_h||_F^2$. Figures [2b](#page-8-1) shows that the ratios between norms approach 1. We then test an unbalanced initialization: $W_1[i, j] \sim N(0, 10^{-4}), W_2[i, j] \sim N(0, 10^{-4})$ and $W_3[i, j] \sim N(0, 10^{-4})$. After 10,000 iterations we have $||W_1||_F^2 = 55.50$, $||W_2||_F^2 = 45.65$ and $||\tilde{W_3}||_F^2 = 45.46$. Figure [2c](#page-8-1) shows that $\left\|W_1\right\|_F^2 - \left\|W_2\right\|_F^2$ and $\left\|W_2\right\|_F^2 - \left\|W_3\right\|_F^2$ are bounded by 9 (and indeed change very little throughout the process), and Figures [2d](#page-8-1) shows that the ratios become close to 1 after about 1,000 iterations.

5 Conclusion and Future Work

In this paper we take a step towards characterizing the invariance imposed by first order algorithms. We show that gradient flow automatically balances the magnitudes of all layers in a deep neural network with homogeneous activations. For the concrete model of asymmetric matrix factorization, we further use the balancedness property to show that gradient descent converges to global minimum. We believe our findings on the invariance in deep models could serve as a fundamental building block for understanding optimization in deep learning. Below we list some future directions.

Other first-order methods. In this paper we focus on the invariance induced by gradient descent. In practice, different acceleration and adaptive methods are also used. A natural future direction is how to characterize the invariance properties of these algorithms.

From gradient flow to gradient descent: a generic analysis? As discussed in Section [3,](#page-6-0) while strong invariance properties hold for gradient flow, in practice one uses gradient descent with positive step sizes and the invariance may only hold approximately because positive step sizes discretize the dynamics. We use specialized techniques for analyzing asymmetric matrix factorization. It would be very interesting to develop a generic approach to analyze the discretization. Recent findings on the connection between optimization and ordinary differential equations [\[Su et al.,](#page-10-18) [2014,](#page-10-18) [Zhang et al.,](#page-10-19) [2018\]](#page-10-19) might be useful for this purpose.

Acknowledgements

We thank Phil Long for his helpful comments on an earlier draft of this paper. JDL acknowledges support from ARO W911NF-11-1-0303.

References

- Pierre-Antoine Absil, Robert Mahony, and Benjamin Andrews. Convergence of the iterates of descent methods for analytic cost functions. *SIAM Journal on Optimization*, 16(2):531–547, 2005.
- Sanjeev Arora, Nadav Cohen, and Elad Hazan. On the optimization of deep networks: Implicit acceleration by overparameterization. *arXiv preprint arXiv:1802.06509*, 2018.
- Peter L Bartlett, David P Helmbold, and Philip M Long. Gradient descent with identity initialization efficiently learns positive definite linear transformations by deep residual networks. *arXiv preprint arXiv:1802.06093*, 2018.
- Alon Brutzkus and Amir Globerson. Globally optimal gradient descent for a convnet with gaussian inputs. *arXiv preprint arXiv:1702.07966*, 2017.
- Alon Brutzkus, Amir Globerson, Eran Malach, and Shai Shalev-Shwartz. Sgd learns overparameterized networks that provably generalize on linearly separable data. *arXiv preprint arXiv:1710.10174*, 2017.
- Anna Choromanska, Mikael Henaff, Michael Mathieu, Gerard Ben Arous, and Yann LeCun. The ´ loss surfaces of multilayer networks. In *Artificial Intelligence and Statistics*, pages 192–204, 2015.
- Francis H Clarke, Yuri S Ledyaev, Ronald J Stern, and Peter R Wolenski. *Nonsmooth analysis and control theory*, volume 178. Springer Science & Business Media, 2008.
- Damek Davis, Dmitriy Drusvyatskiy, Sham Kakade, and Jason D Lee. Stochastic subgradient method converges on tame functions. *arXiv preprint arXiv:1804.07795*, 2018.
- Dmitriy Drusvyatskiy, Alexander D Ioffe, and Adrian S Lewis. Curves of descent. *SIAM Journal on Control and Optimization*, 53(1):114–138, 2015.
- Simon S Du and Jason D Lee. On the power of over-parametrization in neural networks with quadratic activation. *arXiv preprint arXiv:1803.01206*, 2018.
- Simon S Du, Jason D Lee, and Yuandong Tian. When is a convolutional filter easy to learn? *arXiv preprint arXiv:1709.06129*, 2017a.
- Simon S Du, Jason D Lee, Yuandong Tian, Barnabas Poczos, and Aarti Singh. Gradient descent learns one-hidden-layer cnn: Don't be afraid of spurious local minima. *arXiv preprint arXiv:1712.00779*, 2017b.
- C Daniel Freeman and Joan Bruna. Topology and geometry of half-rectified network optimization. *arXiv preprint arXiv:1611.01540*, 2016.
- Rong Ge, Furong Huang, Chi Jin, and Yang Yuan. Escaping from saddle points – online stochastic gradient for tensor decomposition. In *Proceedings of The 28th Conference on Learning Theory*, pages 797–842, 2015.
- Rong Ge, Chi Jin, and Yi Zheng. No spurious local minima in nonconvex low rank problems: A unified geometric analysis. In *Proceedings of the 34th International Conference on Machine Learning*, pages 1233–1242, 2017a.
- Rong Ge, Jason D Lee, and Tengyu Ma. Learning one-hidden-layer neural networks with landscape design. *arXiv preprint arXiv:1711.00501*, 2017b.
- Xavier Glorot and Yoshua Bengio. Understanding the difficulty of training deep feedforward neural networks. In *Proceedings of the thirteenth international conference on artificial intelligence and statistics*, pages 249–256, 2010.
- Benjamin D Haeffele and René Vidal. Global optimality in tensor factorization, deep learning, and beyond. *arXiv preprint arXiv:1506.07540*, 2015.
- Moritz Hardt and Tengyu Ma. Identity matters in deep learning. *arXiv preprint arXiv:1611.04231*, 2016.
- Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep residual learning for image recognition. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 770–778, 2016.
- Kenji Kawaguchi. Deep learning without poor local minima. In *Advances In Neural Information Processing Systems*, pages 586–594, 2016.
- Jason D Lee, Max Simchowitz, Michael I Jordan, and Benjamin Recht. Gradient descent only converges to minimizers. In *Conference on Learning Theory*, pages 1246–1257, 2016.
- Yuanzhi Li and Yang Yuan. Convergence analysis of two-layer neural networks with ReLU activation. *arXiv preprint arXiv:1705.09886*, 2017.
- Siyuan Ma, Raef Bassily, and Mikhail Belkin. The power of interpolation: Understanding the effectiveness of sgd in modern over-parametrized learning. *arXiv preprint arXiv:1712.06559*, 2017.
- Behnam Neyshabur, Ruslan R Salakhutdinov, and Nati Srebro. Path-SGD: Path-normalized optimization in deep neural networks. In *Advances in Neural Information Processing Systems*, pages 2422–2430, 2015a.
- Behnam Neyshabur, Ryota Tomioka, Ruslan Salakhutdinov, and Nathan Srebro. Data-dependent path normalization in neural networks. *arXiv preprint arXiv:1511.06747*, 2015b.
- Quynh Nguyen and Matthias Hein. The loss surface of deep and wide neural networks. *arXiv preprint arXiv:1704.08045*, 2017a.
- Quynh Nguyen and Matthias Hein. The loss surface and expressivity of deep convolutional neural networks. *arXiv preprint arXiv:1710.10928*, 2017b.
- Ioannis Panageas and Georgios Piliouras. Gradient descent only converges to minimizers: Nonisolated critical points and invariant regions. *arXiv preprint arXiv:1605.00405*, 2016.
- Itay Safran and Ohad Shamir. On the quality of the initial basin in overspecified neural networks. In *International Conference on Machine Learning*, pages 774–782, 2016.
- Itay Safran and Ohad Shamir. Spurious local minima are common in two-layer relu neural networks. *arXiv preprint arXiv:1712.08968*, 2017.
- O. Shamir. Are resnets provably better than linear predictors? *arXiv preprint arXiv:1804.06739*, 2018.
- Weijie Su, Stephen Boyd, and Emmanuel Candes. A differential equation for modeling nesterovs accelerated gradient method: Theory and insights. In *Advances in Neural Information Processing Systems*, pages 2510–2518, 2014.
- Yuandong Tian. An analytical formula of population gradient for two-layered ReLU network and its applications in convergence and critical point analysis. *arXiv preprint arXiv:1703.00560*, 2017.
- Stephen Tu, Ross Boczar, Max Simchowitz, Mahdi Soltanolkotabi, and Benjamin Recht. Low-rank solutions of linear matrix equations via procrustes flow. *arXiv preprint arXiv:1507.03566*, 2015.
- Rene Vidal, Joan Bruna, Raja Giryes, and Stefano Soatto. Mathematics of deep learning. *arXiv preprint arXiv:1712.04741*, 2017.
- Jingzhao Zhang, Aryan Mokhtari, Suvrit Sra, and Ali Jadbabaie. Direct runge-kutta discretization achieves acceleration. *arXiv preprint arXiv:1805.00521*, 2018.
- Kai Zhong, Zhao Song, Prateek Jain, Peter L Bartlett, and Inderjit S Dhillon. Recovery guarantees for one-hidden-layer neural networks. *arXiv preprint arXiv:1706.03175*, 2017.
- Pan Zhou and Jiashi Feng. The landscape of deep learning algorithms. *arXiv preprint arXiv:1705.07038*, 2017.

Appendix

A Proofs for Section [2](#page-3-0)

Proof of Theorem [2.2.](#page-4-2) Same as the proof of Theorem [2.1,](#page-4-0) we assume without loss of generality that $L(\boldsymbol{w}) = \ell(f_{\boldsymbol{w}}(\boldsymbol{x}), \boldsymbol{y})$ for some $(\boldsymbol{x}, \boldsymbol{y}) \in \mathbb{R}^d \times \mathbb{R}^p$. We also denote $\boldsymbol{x}^{(h)} = f_{\boldsymbol{w}}^{(h)}(\boldsymbol{x})$ ($\forall h \in [N]$), $\boldsymbol{x}^{(0)} = \boldsymbol{x}$ and $\phi_0(x) = x$.

Now we suppose $\phi_h(x) = x$ for some $h \in [N - 1]$. Denote $u = \phi_{h-1}(x^{(h-1)})$. Then we have $\mathbf{x}^{(h+1)} = \mathbf{W}^{(h+1)}\mathbf{x}^{(h)} = \mathbf{W}^{(h+1)}\mathbf{W}^{(h)}\mathbf{u}$. Using the chain rule, we can directly compute

$$
\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h)}} = \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h)}} \boldsymbol{u}^{\top} = (\boldsymbol{W}^{(h+1)})^{\top} \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}} \boldsymbol{u}^{\top},
$$
\n
$$
\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{W}^{(h+1)}} = \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}} (\boldsymbol{x}^{(h)})^{\top} = \frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}} (\boldsymbol{W}^{(h)} \boldsymbol{u})^{\top}.
$$

Then we have

$$
\frac{\mathrm{d}}{\mathrm{d}t}\left(\boldsymbol{W}^{(h)}(\boldsymbol{W}^{(h)})^{\top}\right) = \boldsymbol{W}^{(h)}\left(\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{W}^{(h)}\right)^{\top} + \left(\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{W}^{(h)}\right)(\boldsymbol{W}^{(h)})^{\top} \n= \boldsymbol{W}^{(h)}\boldsymbol{u}\left(\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}}\right)^{\top}\boldsymbol{W}^{(h+1)} + (\boldsymbol{W}^{(h+1)})^{\top}\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}}\boldsymbol{u}^{\top}(\boldsymbol{W}^{(h)})^{\top},
$$
\n
$$
\frac{\mathrm{d}}{\mathrm{d}t}\left((\boldsymbol{W}^{(h+1)})^{\top}\boldsymbol{W}^{(h+1)}\right) = (\boldsymbol{W}^{(h+1)})^{\top}\left(\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{W}^{(h+1)}\right) + \left(\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{W}^{(h+1)}\right)^{\top}\boldsymbol{W}^{(h+1)} \n= (\boldsymbol{W}^{(h+1)})^{\top}\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}}\boldsymbol{u}^{\top}(\boldsymbol{W}^{(h)})^{\top} + \boldsymbol{W}^{(h)}\boldsymbol{u}\left(\frac{\partial L(\boldsymbol{w})}{\partial \boldsymbol{x}^{(h+1)}}\right)^{\top}\boldsymbol{W}^{(h+1)}.
$$

Comparing the above two equations we know $\frac{d}{dt}$ $\boldsymbol{W}^{(h)} (\boldsymbol{W}^{(h)})^\top - (\boldsymbol{W}^{(h+1)})^\top \boldsymbol{W}^{(h+1)}$ $= 0.$

Proof of Theorem [2.3.](#page-5-3) Same as the proof of Theorem [2.1,](#page-4-0) we assume without loss of generality that $L(\boldsymbol{v}) = L(\boldsymbol{w}) = \ell(f_{\boldsymbol{w}}(\boldsymbol{x}), \boldsymbol{y})$ for $(\boldsymbol{x}, \boldsymbol{y}) \in \mathbb{R}^d \times \mathbb{R}^p$, and denote $\boldsymbol{x}^{(h)} = f_{\boldsymbol{w}}^{(h)}(\boldsymbol{x})$ ($\forall h \in [N]$), $\boldsymbol{x}^{(0)} = \boldsymbol{x}$ and $\phi_0(x) = x$.

Using the chain rule, we have

$$
\frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{v}^{(h+1)}[l]} = \sum_{(k,i):g_{h+1}(k,i)=l} \frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{x}^{(h+1)}[k]} \cdot \phi_h(\boldsymbol{x}^{(h)}[i]), \qquad l \in [d_{h+1}].
$$

Then we have using the sharp chain rule,

$$
\frac{d}{dt} \|\boldsymbol{v}^{(h+1)}\|^2 = 2 \left\langle \boldsymbol{v}^{(h+1)}, \frac{d}{dt} \boldsymbol{v}^{(h+1)} \right\rangle = -2 \left\langle \boldsymbol{v}^{(h+1)}, \frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{v}^{(h+1)}} \right\rangle
$$
\n
$$
= -2 \sum_{l} \sum_{(k,i): g_{h+1}(k,i)=l} \frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{x}^{(h+1)}[k]} \cdot \boldsymbol{v}^{(h+1)}[l] \cdot \phi_h(\boldsymbol{x}^{(h)}[i])
$$
\n
$$
= -2 \sum_{(k,i)} \frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{x}^{(h+1)}[k]} \cdot \boldsymbol{W}^{(h+1)}[k, i] \cdot \phi_h(\boldsymbol{x}^{(h)}[i])
$$
\n
$$
= -2 \sum_{k} \frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{x}^{(h+1)}[k]} \cdot \boldsymbol{x}^{(h+1)}[k]
$$
\n
$$
= -2 \left\langle \frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{x}^{(h+1)}}, \boldsymbol{x}^{(h+1)} \right\rangle.
$$
\n(10)

Substituting h with $h-1$ in [\(10\)](#page-12-1) gives $\frac{d}{dt}||v^{(h)}||^2 = -2$ $\partial L(\bm{v})$ $\frac{\partial L(\boldsymbol{v})}{\partial \boldsymbol{x}^{(h)}}, \boldsymbol{x}^{(h)}$, which further implies

$$
\frac{d}{dt} \|\mathbf{v}^{(h)}\|^2 = -2 \left\langle \frac{\partial L(\mathbf{v})}{\partial \mathbf{x}^{(h)}}, \mathbf{x}^{(h)} \right\rangle = -2 \sum_i \frac{\partial L(\mathbf{v})}{\partial \mathbf{x}^{(h)}[i]} \cdot \mathbf{x}^{(h)}[i]
$$
\n
$$
= -2 \sum_i \sum_k \frac{\partial L(\mathbf{v})}{\partial \mathbf{x}^{(h+1)}[k]} \cdot \mathbf{W}^{(h+1)}[k, i] \cdot \phi'_h(\mathbf{x}^{(h)}[i]) \cdot \mathbf{x}^{(h)}[i]
$$
\n
$$
= -2 \sum_k \frac{\partial L(\mathbf{v})}{\partial \mathbf{x}^{(h+1)}[k]} \sum_i \mathbf{W}^{(h+1)}[k, i] \cdot \phi_h(\mathbf{x}^{(h)}[i])
$$
\n
$$
= -2 \sum_k \frac{\partial L(\mathbf{v})}{\partial \mathbf{x}^{(h+1)}[k]} \cdot \mathbf{x}^{(h+1)}[k]
$$
\n
$$
= -2 \left\langle \frac{\partial L(\mathbf{v})}{\partial \mathbf{x}^{(h+1)}}, \mathbf{x}^{(h+1)} \right\rangle.
$$
\n(11)

The proof is finished by combining (10) and (11) .

 \Box

B Proof for Rank- r Matrix Factorization (Theorem [3.1\)](#page-6-5)

In this section we give the full proof of Theorem [3.1.](#page-6-5)

First we recall the gradient of our objective function $f(U, V) = \frac{1}{2}$ $\left\Vert \boldsymbol{U}\boldsymbol{V}^{\top}-\boldsymbol{M}^{*}\right\Vert _{F}^{2}$ $\frac{2}{F}$:

$$
\frac{\partial f(\boldsymbol{U},\boldsymbol{V})}{\partial \boldsymbol{U}}=(\boldsymbol{U}\boldsymbol{V}^{\top}-\boldsymbol{M}^{*})\boldsymbol{V},\qquad \frac{\partial f(\boldsymbol{U},\boldsymbol{V})}{\partial \boldsymbol{V}}=(\boldsymbol{U}\boldsymbol{V}^{\top}-\boldsymbol{M}^{*})^{\top}\boldsymbol{U}.
$$

We also need to calculate the Hessian $\nabla^2 f(U, V)$. The Hessian can be viewed as a matrix that We also need to calculate the Hessian $\nabla^2 f(U, V)$. The Hessian can be viewed as a r operates on vectorized matrices of dimension $(d_1 + d_2) \times r$ (i.e., the same shape as $\begin{pmatrix} U \\ V \end{pmatrix}$ $\begin{bmatrix} V \\ V \end{bmatrix}$). Then, for any $W \in \mathbb{R}^{(d_1+d_2)\times r}$, the Hessian $\nabla^2 f(\boldsymbol{W})$ defines a quadratic form

$$
[\nabla^2 f(\boldsymbol{W})](\boldsymbol{A},\boldsymbol{B}) = \sum_{i,j,k,l} \frac{\partial^2 f(\boldsymbol{W})}{\partial \boldsymbol{W}[i,j] \partial \boldsymbol{W}[k,l]} \boldsymbol{A}[i,j] \boldsymbol{B}[k,l], \qquad \forall \boldsymbol{A}, \boldsymbol{B} \in \mathbb{R}^{(d_1+d_2) \times r}.
$$

With this notation, we can express the Hessian $\nabla^2 f(U, V)$ as follows:

$$
[\nabla^2 f(U, V)](\Delta, \Delta) = 2 \langle UV^\top - M^*, \Delta_U \Delta_V^\top \rangle + \|U \Delta_V^\top + \Delta_U V^\top\|_F^2, \forall \Delta = \left(\frac{\Delta_U}{\Delta_V}\right), \Delta_U \in \mathbb{R}^{d_1 \times r}, \Delta_V \in \mathbb{R}^{d_2 \times r}.
$$
\n(12)

Now we use the expression of the Hessian to prove that $f(U, V)$ is locally smooth when both arguments U and V are bounded.

Lemma B.1 (Smoothness over a bounded set). *For any* $c > 0$, constrained on the set $S =$ $\{(U,V) : U \in \mathbb{R}^{d_1 \times r}, V \in \mathbb{R}^{d_2 \times r}, \|U\|_F^2 \leqslant c\|M^*\|_F, \|V\|_F^2 \leqslant c\|M^*\|_F\}$, the function f is $\left(\left(6c+2\right)\|M^*\|_F\right)$ -smooth.

Proof. We prove smoothness by giving an upper bound on $\lambda_{\max}(\nabla^2 f(U, V))$ for any $(U, V) \in S$. For any $(U, V) \in S$ and any $\Delta = \begin{pmatrix} \Delta_U \\ \Delta_U \end{pmatrix}$ $\left(\frac{\Delta U}{\Delta V}\right)$ $(\Delta_U \in \mathbb{R}^{d_1 \times r}, \Delta_V \in \mathbb{R}^{d_2 \times r})$, from [\(12\)](#page-13-2) we have

$$
\begin{split} &\left[\nabla^2 f(\boldsymbol{U},\boldsymbol{V})\right] (\boldsymbol{\Delta},\boldsymbol{\Delta}) \\ &\leqslant 2\left\|\boldsymbol{U}\boldsymbol{V}^\top - \boldsymbol{M}^*\right\|_F \left\|\boldsymbol{\Delta}_U\boldsymbol{\Delta}_\boldsymbol{V}^\top\right\|_F + \left\|\boldsymbol{U}\boldsymbol{\Delta}_V^\top + \boldsymbol{\Delta}_U\boldsymbol{V}^\top\right\|_F^2 \\ &\leqslant 2\left(\left\|\boldsymbol{U}\right\|_F \left\|\boldsymbol{V}^\top\right\|_F + \left\|\boldsymbol{M}^*\right\|_F\right) \left\|\boldsymbol{\Delta}_U\right\|_F \left\|\boldsymbol{\Delta}_V^\top\right\|_F + \left(\left\|\boldsymbol{U}\right\|_F \left\|\boldsymbol{\Delta}_V^\top\right\|_F + \left\|\boldsymbol{\Delta}_U\right\|_F \left\|\boldsymbol{V}^\top\right\|_F\right)^2 \end{split}
$$

$$
\leq 2 (c \| {\boldsymbol M}^* \|_F + \| {\boldsymbol M}^* \|_F) \| {\boldsymbol \Delta} \|^2_F + \left(2 \sqrt{c \| {\boldsymbol M}^* \|_F} \cdot \| {\boldsymbol \Delta} \|_F \right)^2
$$

= $(6c+2) \| {\boldsymbol M}^* \|_F \| {\boldsymbol \Delta} \|^2_F.$

This implies $\lambda_{\max}(\nabla^2 f({\boldsymbol U},{\boldsymbol V})) \leqslant (6c+2) \left\| {\boldsymbol M}^* \right\|_F$.

B.1 Proof of Lemma [3.1](#page-6-6)

Recall the following three properties we want to prove in Lemma [3.1,](#page-6-6) which we call $A(t)$, $B(t)$ and $C(t)$, respectively:

$$
\mathcal{A}(t): \qquad \left\| \boldsymbol{U}_t^{\top} \boldsymbol{U}_t - \boldsymbol{V}_t^{\top} \boldsymbol{V}_t \right\|_F \leq \epsilon, \n\mathcal{B}(t): \qquad f(\boldsymbol{U}_t, \boldsymbol{V}_t) \leq f(\boldsymbol{U}_{t-1}, \boldsymbol{V}_{t-1}) \leq \cdots \leq f(\boldsymbol{U}_0, \boldsymbol{V}_0) \leq 2 \left\| \boldsymbol{M}^* \right\|_F^2, \n\mathcal{C}(t): \qquad \left\| \boldsymbol{U}_t \right\|_F^2 \leq \frac{5\sqrt{r}}{|\boldsymbol{M}^*|_F, \left\| \boldsymbol{V}_t \right\|_F^2} \leq \frac{5\sqrt{r}}{|\boldsymbol{M}^*|_F}.
$$

We use induction to prove these statements. For $t = 0$, we can make the Gaussian variance in the initialization sufficiently small such that with high probability we have

$$
\left\|\boldsymbol{U}_0\right\|_F^2 \leqslant \epsilon, \qquad \left\|\boldsymbol{V}_0\right\|_F^2 \leqslant \epsilon, \qquad \left\|\boldsymbol{U}_0^\top \boldsymbol{U}_0 - \boldsymbol{V}_0^\top \boldsymbol{V}_0\right\|_F \leqslant \frac{\epsilon}{2}
$$

From now on we assume they are all satisfied. Then $A(0)$ is already satisfied, $C(0)$ is satisfied because ϵ $\|M^*\|_F$, and $\mathcal{B}(0)$ can be verified by $f(U_0, V_0) = \frac{1}{2} ||U_0 V_0^\top - M^*||_F^2$ fied because $\epsilon < \|M^*\|_F$, and $\mathcal{B}(0)$ can be verified by $f(U_0, V_0) = \frac{1}{2} \|U_0 V_0^{\dagger} - M^*\|_F^2 \le \|U_0 V_0^{\dagger}\|_F^2 + \|M^*\|_F^2 \le \|U_0\|_F^2 + \|M^*\|_F^2 + \|M^*\|_F^2 \le \epsilon^2 + \|M^*\|_F^2 \le 2\|M^*\|_F^2$. $\frac{a}{\parallel^2}$ $\frac{2}{F}+\left\Vert \boldsymbol{M}^{\ast}\right\Vert _{F}^{2}\leqslant\left\Vert \boldsymbol{U}_{0}\right\Vert _{F}^{2}$ $\begin{array}{c} \mathbf{v} \mathbf{v} \mathbf{v} \ \mathbf{v} \mathbf{v} \end{array}$ י
|² $\frac{2}{F} + \left\| \bm{M^*} \right\|_F^2 \leqslant \epsilon^2 + \left\| \bm{M^*} \right\|_F^2 \leqslant 2 \left\| \bm{M^*} \right\|_F^2.$

To prove $A(t)$, $B(t)$ and $C(t)$ for all t, we prove the following three claims. Since we have $A(0)$, $B(0)$ and $C(0)$, if the following claims are all true, the proof will be completed by induction.

(i) $\mathcal{B}(0), \ldots, \mathcal{B}(t), \mathcal{C}(0), \ldots, \mathcal{C}(t) \implies \mathcal{A}(t + 1);$ (ii) $\mathcal{B}(0), \ldots, \mathcal{B}(t), \mathcal{C}(t) \implies \mathcal{B}(t + 1);$ (iii) $\mathcal{A}(t), \mathcal{B}(t) \implies \mathcal{C}(t)$. Claim B.1. $\mathcal{B}(0), \ldots, \mathcal{B}(t), \mathcal{C}(0), \ldots, \mathcal{C}(t) \implies \mathcal{A}(t+1)$.

Proof. Using the update rule [\(9\)](#page-6-3) we can calculate

$$
\begin{aligned} &\bm{U}_{t+1}^{\top} \bm{U}_{t+1} - \bm{V}_{t+1}^{\top} \bm{V}_{t+1} \\& = \left(\bm{U}_{t} - \eta_{t} (\bm{U}_{t} \bm{V}_{t}^{\top} - \bm{M}^{*}) \bm{V}_{t} \right)^{\top} \left(\bm{U}_{t} - \eta_{t} (\bm{U}_{t} \bm{V}_{t}^{\top} - \bm{M}^{*}) \bm{V}_{t} \right) \\& - \left(\bm{V}_{t} - \eta_{t} (\bm{U}_{t} \bm{V}_{t}^{\top} - \bm{M}^{*})^{\top} \bm{U}_{t} \right)^{\top} \left(\bm{V}_{t} - \eta_{t} (\bm{U}_{t} \bm{V}_{t}^{\top} - \bm{M}^{*})^{\top} \bm{U}_{t} \right) \\& = \bm{U}_{t}^{\top} \bm{U}_{t} - \bm{V}_{t}^{\top} \bm{V}_{t} + \eta_{t}^{2} \left(\bm{V}_{t}^{\top} \bm{R}_{t}^{\top} \bm{R}_{t} \bm{V}_{t} - \bm{U}_{t}^{\top} \bm{R}_{t}^{\top} \bm{R}_{t} \bm{U}_{t} \right), \end{aligned}
$$

where $\boldsymbol{R}_t = \boldsymbol{U}_t \boldsymbol{V}_t^\top - \boldsymbol{M^*}.$ Then we have

$$
\|U_{t+1}^{\top}U_{t+1} - V_{t+1}^{\top}V_{t+1}\|_{F} \n\leq \|U_{t}^{\top}U_{t} - V_{t}^{\top}V_{t}\|_{F} + \eta_{t}^{2} \left(\|V_{t}^{\top}R_{t}^{\top}R_{t}V_{t}\|_{F} + \|U_{t}^{\top}R_{t}^{\top}R_{t}U_{t}\|_{F}\right) \n\leq \|U_{t}^{\top}U_{t} - V_{t}^{\top}V_{t}\|_{F} + \eta_{t}^{2} \left(\|V_{t}\|_{F}^{2} \|R_{t}\|_{F}^{2} + \|U_{t}\|_{F}^{2} \|R_{t}\|_{F}^{2}\right) \n= \|U_{t}^{\top}U_{t} - V_{t}^{\top}V_{t}\|_{F} + 2\eta_{t}^{2} \left(\|V_{t}\|_{F}^{2} + \|U_{t}\|_{F}^{2}\right) f(U_{t}, V_{t}) \n\leq \|U_{t}^{\top}U_{t} - V_{t}^{\top}V_{t}\|_{F} + 2\eta_{t}^{2} \cdot 10\sqrt{r} \|M^{*}\|_{F} \cdot 2 \|M^{*}\|_{F}^{2},
$$
\n(13)

where the last line is due to $\mathcal{B}(t)$ and $\mathcal{C}(t)$.

Since we have $\mathcal{B}(t')$ and $\mathcal{C}(t')$ for all $t' \leq t$, [\(13\)](#page-14-0) is still true when substituting t with any $t' \leq t$. Summing all of them and noting $\left\|U_0^\top U_0 - V_0^\top V_0\right\|_F \leq \frac{\epsilon}{2}$, we get \mathbb{R} is the set of \mathbb{R} in the set of \mathbb{R}

$$
\big\| \bm{U}_{t+1}^{\top} \bm{U}_{t+1} - \bm{V}_{t+1}^{\top} \bm{V}_{t+1} \big\|_F
$$

 \Box

.

$$
\leq \left\| \boldsymbol{U}_0^{\top} \boldsymbol{U}_0 - \boldsymbol{V}_0^{\top} \boldsymbol{V}_0 \right\|_F + 40\sqrt{r} \left\| \boldsymbol{M}^* \right\|_F^3 \sum_{i=0}^t \eta_i^2
$$

$$
\leq \frac{\epsilon}{2} + 40\sqrt{r} \left\| \boldsymbol{M}^* \right\|_F^3 \sum_{i=0}^t \frac{1}{(i+1)^2} \cdot \frac{\epsilon/r}{100^2 \left\| \boldsymbol{M}^* \right\|_F^3}
$$

$$
\leq \epsilon.
$$

Therefore we have proved $A(t + 1)$.

Claim B.2. $\mathcal{B}(0), \ldots, \mathcal{B}(t), \mathcal{C}(t) \implies \mathcal{B}(t + 1)$.

Proof. Note that we only need to show $f(U_{t+1},V_{t+1}) \leq f(U_t,V_t)$. We prove this using the standard analysis of gradient descent, for which we need the smoothness of the objective function f (Lemma [B.1\)](#page-13-3). We first need to bound $||U_t||_F$, $||V_t||_F$, $||U_{t+1}||_F$ and $||V_{t+1}||_F$. We know from $\mathcal{C}(t)$ that $||U_t||_F^2 \le 5\sqrt{r}||M^*||_F$ and $||V_t||_F^2 \le 5\sqrt{r}||M^*||_F$. We can also bound $||U_{t+1}||_F^2$ and $||V_{t+1}||_F^2$ easily from the GD update rule:

$$
\|U_{t+1}\|_{F}^{2}
$$
\n
$$
= \|U_{t} - \eta_{t}(U_{t}V_{t}^{\top} - M^{*})V_{t}\|_{F}^{2}
$$
\n
$$
\leq 2\|U_{t}\|_{F}^{2} + 2\eta_{t}^{2}\|U_{t}V_{t}^{\top} - M^{*}\|_{F}^{2}\|V_{t}\|_{F}^{2}
$$
\n
$$
\leq 2\cdot 5\sqrt{r}\|M^{*}\|_{F} + 2\eta_{t}^{2}\cdot 2f(U_{t}, V_{t})\cdot 5\sqrt{r}\|M^{*}\|_{F}
$$
\n
$$
\leq 10\sqrt{r}\|M^{*}\|_{F} + 2\cdot \frac{\epsilon/r}{100^{2}(t+1)^{2}\|M^{*}\|_{F}^{3}} \cdot 4\|M^{*}\|_{F}^{2}\cdot 5\sqrt{r}\|M^{*}\|_{F}
$$
\n
$$
\leq 10\sqrt{r}\|M^{*}\|_{F} + \frac{\epsilon}{100}
$$
\n
$$
\leq 11\sqrt{r}\|M^{*}\|_{F} + \frac{\epsilon}{100}
$$
\n
$$
\leq 11\sqrt{r}\|M^{*}\|_{F}.
$$
\n(using $\epsilon < \|M^{*}\|_{F})$

Let $\beta = (66\sqrt{r} + 2) \|M^*\|_F$. From Lemma [B.1,](#page-13-3) f is β -smooth over $S = \{(\boldsymbol{U}, \boldsymbol{V}) : \|\boldsymbol{U}\|_F^2 \le 11\sqrt{r} \|\boldsymbol{M}^*\|_F$, $\|\boldsymbol{V}\|_F^2 \le 11\sqrt{r} \|\boldsymbol{M}^*\|_F$. Also note that $\eta_t < \frac{1}{\beta}$ by our choice. Then using smoo ness we have $f(Tt)$ T

$$
f(U_{t+1}, V_{t+1})
$$
\n
$$
\leq f(U_t, V_t) + \left\langle \nabla f(U_t, V_t), \left(\frac{U_{t+1}}{V_{t+1}} \right) - \left(\frac{U_t}{V_t} \right) \right\rangle + \frac{\beta}{2} \left\| \left(\frac{U_{t+1}}{V_{t+1}} \right) - \left(\frac{U_t}{V_t} \right) \right\|_F^2
$$
\n
$$
= f(U_t, V_t) - \eta_t \left\| \nabla f(U_t, V_t) \right\|_F^2 + \frac{\beta}{2} \eta_t^2 \left\| \nabla f(U_t, V_t) \right\|_F^2
$$
\n
$$
\leq f(U_t, V_t) - \frac{\eta_t}{2} \left\| \nabla f(U_t, V_t) \right\|_F^2.
$$
\n(14)

Therefore we have shown $\mathcal{B}(t + 1)$.

Claim B.3. $\mathcal{A}(t), \mathcal{B}(t) \implies \mathcal{C}(t)$.

Proof. From $\mathcal{B}(t)$ we know $\frac{1}{2}$ $\left\Vert \boldsymbol{U}_{t}\boldsymbol{V}_{t}^{\top}-\boldsymbol{M}^{\ast}\right\Vert$ \mathbb{I}^2 $\frac{2}{F} \leq 2 \left\| \boldsymbol{M}^* \right\|_F^2$ which implies $\begin{aligned} \big\| \boldsymbol{U}_t \boldsymbol{V}_t^\top \end{aligned}$ $\Big\|_F \leq$ $3\left\Vert \mathbf{\mathit{M}}^{\ast}\right\Vert _{F}.$ Therefore it suffices to prove

$$
\|UV^{\top}\|_{F} \leq 3\|M^{*}\|_{F},\|U^{\top}U-V^{\top}V\|_{F} \leq \epsilon \implies \|U\|_{F}^{2} \leq 5\sqrt{r}\|M^{*}\|_{F},\|V\|_{F}^{2} \leq 5\sqrt{r}\|M^{*}\|_{F}.
$$
\n(15)

Now we prove [\(15\)](#page-15-0). Consider the SVD $U = \Phi \Sigma \Psi^{\top}$, where $\Phi \in \mathbb{R}^{d_1 \times d_1}$ and $\Psi \in \mathbb{R}^{r \times r}$ are orthogonal matrices, and $\Sigma \in \mathbb{R}^{d_1 \times r}$ is a diagonal matrix. Let $\sigma_i = \Sigma[i, i]$ $(i \in [r])$ which are all the singular values of U. Define $\tilde{V} = V \Psi$. Then we have

$$
3\left\| \boldsymbol{M}^* \right\|_F \geqslant \left\| \boldsymbol{U} \boldsymbol{V}^\top \right\|_F = \left\| \boldsymbol{\Phi} \boldsymbol{\Sigma} \boldsymbol{\Psi}^\top \boldsymbol{\Psi} \boldsymbol{\tilde{V}}^\top \right\|_F = \left\| \boldsymbol{\Sigma} \boldsymbol{\tilde{V}}^\top \right\|_F = \sqrt{\sum_{i=1}^r \sigma_i^2 \left\| \boldsymbol{\tilde{V}}[:,i] \right\|^2}
$$

 \Box

 \Box

and

$$
\begin{aligned} &\epsilon \geqslant \left\| \boldsymbol{U}^\top \boldsymbol{U} - \boldsymbol{V}^\top \boldsymbol{V} \right\|_F = \left\| \boldsymbol{\Psi} \boldsymbol{\Sigma}^\top \boldsymbol{\Phi}^\top \boldsymbol{\Phi} \boldsymbol{\Sigma} \boldsymbol{\Psi}^\top - \boldsymbol{\Psi} \widetilde{\boldsymbol{V}}^\top \widetilde{\boldsymbol{V}} \boldsymbol{\Psi}^\top \right\|_F = \left\| \boldsymbol{\Sigma}^\top \boldsymbol{\Sigma} - \widetilde{\boldsymbol{V}}^\top \widetilde{\boldsymbol{V}} \right\|_F \\ &\geqslant \sqrt{\sum_{i=1}^r \left(\sigma_i^2 - \left\| \widetilde{\boldsymbol{V}}[:,i] \right\|^2 \right)^2}. \end{aligned}
$$

Using the above two inequalities we get

$$
\sum_{i=1}^{r} \sigma_i^4 \le \sum_{i=1}^{r} \left(\sigma_i^4 + \left\| \tilde{V}[:,i] \right\|^4 \right) = \sum_{i=1}^{r} \left(\sigma_i^2 - \left\| \tilde{V}[:,i] \right\|^2 \right)^2 + 2 \sum_{i=1}^{r} \sigma_i^2 \left\| \tilde{V}[:,i] \right\|^2
$$

$$
\le \epsilon^2 + 2 \left(3 \left\| M^* \right\|_F \right)^2 \le 19 \left\| M^* \right\|_F^2.
$$

Then by the Cauchy-Schwarz inequality we have

$$
||U||_F^2 = \sum_{i=1}^r \sigma_i^2 \le \sqrt{r \sum_{i=1}^r \sigma_i^4} \le \sqrt{r \cdot 19 ||M^*||_F^2} \le 5\sqrt{r} ||M^*||_F^2.
$$

Similarly, we also have $\|{\bm{V}}\|_F^2\leqslant 5\sqrt{r}\,\|{\bm{M}}^*\|_F.$ Therefore we have proved [\(15\)](#page-15-0).

\Box

 \Box

B.2 Convergence to a Stationary Point

With the balancedness and boundedness properties in Lemma [3.1,](#page-6-6) it is then standard to show that (U_t, V_t) converges to a stationary point of f.

Lemma B.2. Under the setting of Theorem [3.1,](#page-6-5) with high probability $\lim_{t\to\infty} (U_t, V_t) = (\bar{U}, \bar{V})$ **exists, and** (\bar{U}, \bar{V}) is a stationary point of f. Furthermore, (\bar{U}, \bar{V}) satisfies $\|\bar{U}^{\top}\bar{U} - \bar{V}^{\top}\bar{V}\| \leq \epsilon$.

Proof. We assume the three properties in Lemma [3.1](#page-6-6) hold, which happens with high probability. Then from [\(14\)](#page-15-1) we have

$$
f(\boldsymbol{U}_{t+1}, \boldsymbol{V}_{t+1}) \leq f(\boldsymbol{U}_t, \boldsymbol{V}_t) - \frac{\eta_t}{2} \|\nabla f(\boldsymbol{U}_t, \boldsymbol{V}_t)\|_F^2
$$

= $f(\boldsymbol{U}_t, \boldsymbol{V}_t) - \frac{1}{2} \|\nabla f(\boldsymbol{U}_t, \boldsymbol{V}_t)\|_F \left\| \begin{pmatrix} \boldsymbol{U}_{t+1} \\ \boldsymbol{V}_{t+1} \end{pmatrix} - \begin{pmatrix} \boldsymbol{U}_t \\ \boldsymbol{V}_t \end{pmatrix} \right\|_F.$ (16)

Under the above descent condition, the result of [Absil et al.](#page-9-16) [\[2005\]](#page-9-16) says that the iterates either diverge to infinity or converge to a fixed point. According to Lemma [3.1,](#page-6-6) $\{(U_t, V_t)\}_{t=1}^{\infty}$ are all bounded, so they have to converge to a fixed point (U, V) as $t \to \infty$.

Next, from [\(16\)](#page-16-1) we know that $\sum_{t=1}^{\infty} \frac{\eta_t}{2} \|\nabla f(\boldsymbol{U}_t, \boldsymbol{V}_t)\|_F^2 \leq f(\boldsymbol{U}_0, \boldsymbol{V}_0)$ is bounded. Notice that η_t scales like 1/t. So we must have $\liminf_{t\to\infty} \|\nabla f(\boldsymbol{U}_t,\boldsymbol{V}_t)\|_{F} = 0$. Then according to the smooth-ness of f in a bounded region (Lemma [B.1\)](#page-13-3) we conclude $\nabla f(\bar{U}, \bar{V}) = 0$, i.e., (\bar{U}, \bar{V}) is a stationary point.

The second part of the lemma is evident according to Lemma [3.1](#page-6-6) (i).

B.3 Proof of Lemma [3.2](#page-7-1)

The main idea in the proof is similar to [Ge et al.](#page-9-0) [\[2017a\]](#page-9-0). We want to find a direction Δ such that either $[\nabla^2 f(U,V)](\Delta, \Delta)$ is negative or (U,V) is close to a global minimum. We show that this is possible when $[V][(\Delta,\Delta) \text{ is not} \brack \|U^\top U - V^\top V\|$ gative o $\Big\|_F \leqslant \epsilon.$

First we define some notation. Take the SVD $M^* = \Phi^* \Sigma^* \Psi^{*\top}$, where $\Phi^* \in \mathbb{R}^{d_1 \times r}$ and $\Psi^* \in \mathbb{R}^{d_2 \times r}$ have orthonormal columns and $\Sigma^* \in \mathbb{R}^{r \times r}$ is diagonal. Denote $U^* = \Phi^* (\Sigma^*)^{1/2}$ and $V^* = \Psi^*(\Sigma^*)^{1/2}$. Then we have $U^*V^{*\top} = M^*$ (i.e., (U^*,V^*) is a global minimum) and $\boldsymbol{U}^{*\top}\boldsymbol{U}^* = \boldsymbol{V}^{*\top}\boldsymbol{V}^*.$

Let
$$
M = UV^{\top}
$$
, $W = \begin{pmatrix} U \\ V \end{pmatrix}$ and $W^* = \begin{pmatrix} U^* \\ V^* \end{pmatrix}$. Define
\n
$$
R = \operatorname{argmin}_{R' \in \mathbb{R}^{r \times r}, \text{ orthogonal}} \|W - W^* R'\|_F
$$

$$
\Delta = W - W^* R.
$$

We will show that Δ is the desired direction. Recall [\(12\)](#page-13-2):

$$
[\nabla^2 f(\boldsymbol{U}, \boldsymbol{V})](\boldsymbol{\Delta}, \boldsymbol{\Delta}) = 2 \langle \boldsymbol{M} - \boldsymbol{M}^*, \boldsymbol{\Delta}_{\boldsymbol{U}} \boldsymbol{\Delta}_{\boldsymbol{V}}^\top \rangle + \left\| \boldsymbol{U} \boldsymbol{\Delta}_{\boldsymbol{V}}^\top + \boldsymbol{\Delta}_{\boldsymbol{U}} \boldsymbol{V}^\top \right\|_F^2, \tag{17}
$$

where $\Delta =$ Δ_U Δ_{V} , $\Delta_U \in \mathbb{R}^{d_1 \times r}$, $\Delta_V \in \mathbb{R}^{d_2 \times r}$. We consider the two terms in [\(17\)](#page-17-0) separately.

For the first term in (17) , we have:

Claim B.4. $\langle M - M^*, \Delta_U \Delta_V^{\dagger} \rangle$ $=-\left\Vert \boldsymbol{M}-\boldsymbol{M}^{*}\right\Vert _{F}^{2}.$

Proof. Since (U, V) is a stationary point of f, we have the first-order optimality condition:

$$
\frac{\partial f(\boldsymbol{U}, \boldsymbol{V})}{\partial \boldsymbol{U}} = (\boldsymbol{M} - \boldsymbol{M}^*)\boldsymbol{V} = \boldsymbol{0}, \qquad \frac{\partial f(\boldsymbol{U}, \boldsymbol{V})}{\partial \boldsymbol{V}} = (\boldsymbol{M} - \boldsymbol{M}^*)^{\top} \boldsymbol{U} = \boldsymbol{0}.
$$
 (18)

Note that $\Delta_U = U - U^*R$ and $\Delta_V = V - V^*R$. We have

$$
\begin{aligned} &\langle \boldsymbol{M}-\boldsymbol{M}^*,\boldsymbol{\Delta_U}\boldsymbol{\Delta_V^\top}\rangle \\ &=\big\langle \boldsymbol{M}-\boldsymbol{M}^*, (\boldsymbol{U}-\boldsymbol{U}^*\boldsymbol{R})(\boldsymbol{V}-\boldsymbol{V}^*\boldsymbol{R})^\top \big\rangle \\ &=\big\langle \boldsymbol{M}-\boldsymbol{M}^*,\boldsymbol{M}-\boldsymbol{U}^*\boldsymbol{R}\boldsymbol{V}^\top-\boldsymbol{U}\boldsymbol{R}^\top\boldsymbol{V}^{*\top}+\boldsymbol{M}^*\big\rangle \\ &=\langle \boldsymbol{M}-\boldsymbol{M}^*,\boldsymbol{M}^*\rangle \\ &=\langle \boldsymbol{M}-\boldsymbol{M}^*,\boldsymbol{M}^*-\boldsymbol{M}\rangle \\ &=-\|\boldsymbol{M}-\boldsymbol{M}^*\|_F^2\,, \end{aligned}
$$

where we have used the following consequences of [\(18\)](#page-17-1):

$$
\langle \mathbf{M} - \mathbf{M}^*, \mathbf{M} \rangle = \langle \mathbf{M} - \mathbf{M}^*, \mathbf{U}\mathbf{V}^\top \rangle = 0,
$$

$$
\langle \mathbf{M} - \mathbf{M}^*, \mathbf{U}^* \mathbf{R} \mathbf{V}^\top \rangle = 0,
$$

$$
\langle \mathbf{M} - \mathbf{M}^*, \mathbf{U} \mathbf{R}^\top \mathbf{V}^{*\top} \rangle = 0.
$$

 \Box

The second term in (17) has the following upper bound:

Claim B.5. $||U\Delta_V + \Delta_U V||_F^2 \le ||M - M^*||_F^2 + \frac{1}{2}\epsilon^2$.

Proof. We make use of the following identities, all of which can be directly verified by plugging in definitions:

$$
U\Delta_V^{\top} + \Delta_U V^{\top} = \Delta_U \Delta_V^{\top} + M - M^*,\tag{19}
$$

$$
\left\| \Delta \Delta^{\top} \right\|_{F}^{2} = 4 \left\| \Delta_{U} \Delta_{V}^{\top} \right\|_{F}^{2} + \left\| \Delta_{U}^{\top} \Delta_{U} - \Delta_{V}^{\top} \Delta_{V} \right\|_{F}^{2},\tag{20}
$$

$$
\|WW^{\top} - W^*W^{*\top}\|_F^2 = 4\|M - M^*\|_F^2 - 2\|U^{\top}U^* - V^{\top}V^*\|_F^2 + \|U^{\top}U - V^{\top}V\|_F^2 + \|U^*U^* - V^{*\top}V^*\|_F^2.
$$
\n(21)

We also need the following inequality, which is [\[Ge et al.,](#page-9-0) [2017a,](#page-9-0) Lemma 6]:

$$
\left\| \Delta \Delta^{\top} \right\|_{F}^{2} \leqslant 2 \left\| WW^{\top} - W^* W^{* \top} \right\|_{F}^{2}.
$$
 (22)

Now we can prove the desired bound as follows:

$$
\|U\Delta_V + \Delta_U V\|_F^2
$$

= $\|\Delta_U \Delta_V^{\top} + M - M^*\|_F^2$
= $\|\Delta_U \Delta_V^{\top}\|_F^2 + 2\langle M - M^*, \Delta_U \Delta_V^{\top}\rangle + \|M - M^*\|_F^2$ ((19))

and

$$
= \left\| \Delta_U \Delta_V^{\top} \right\|_F^2 - \left\| M - M^* \right\|_F^2 \tag{Claim B.4}
$$

$$
\leqslant \frac{1}{4} \left\| \Delta \Delta^{\top} \right\|_{F}^{2} - \left\| M - M^* \right\|_{F}^{2} \tag{20}
$$

$$
\leqslant \frac{1}{2} \left\| \boldsymbol{W} \boldsymbol{W}^{\top} - \boldsymbol{W}^* \boldsymbol{W}^{* \top} \right\|_{F}^{2} - \left\| \boldsymbol{M} - \boldsymbol{M}^* \right\|_{F}^{2}
$$
\n(22)

$$
= 2 \left\| M - M^* \right\|_F^2 - \left\| U^\top U^* - V^\top V^* \right\|_F^2 + \frac{1}{2} \left\| U^\top U - V^\top V \right\|_F^2 + \frac{1}{2} \left\| U^{* \top} U^* - V^{* \top} V^* \right\|_F^2 - \left\| M - M^* \right\|_F^2 \leqslant \left\| M - M^* \right\|_F^2 + \frac{1}{2} \epsilon^2,
$$
\n(21)

where in the last line we have used $U^{*T}U^* = V^{*T}V^*$ and $\|\boldsymbol{U}^\top \boldsymbol{U} - \boldsymbol{V}^\top \boldsymbol{V}\|$ $\|\leqslant \epsilon.$

 \Box

Using Claims $B.4$ and $B.5$, we obtain an upper bound on (17) :

$$
[\nabla^2 f(\boldsymbol{U}, \boldsymbol{V})](\boldsymbol{\Delta}, \boldsymbol{\Delta}) \leq -\|\boldsymbol{M} - \boldsymbol{M}^*\|_F^2 + \frac{1}{2}\epsilon^2.
$$

Therefore, we have either $\left\| \boldsymbol{U} \boldsymbol{V}^\top - \boldsymbol{M}^* \right\|_F \; = \; \left\| \boldsymbol{M} - \boldsymbol{M}^* \right\|_F \; \leqslant \; \epsilon \, \, \text{or} \, \, [\nabla^2 f(\boldsymbol{U}, \boldsymbol{V})](\boldsymbol{\Delta}, \boldsymbol{\Delta}) \; \leqslant$ $-\frac{1}{2}\epsilon^2$ < 0. In the latter case, (U, V) is a strict saddle point of f. This completes the proof of Lemma [3.2.](#page-7-1)

B.4 Finishing the Proof of Theorem [3.1](#page-6-5)

Theorem [3.1](#page-6-5) is a direct corollary of Lemma [B.2,](#page-16-0) Lemma [3.2,](#page-7-1) and the fact that gradient descent does not converge to a strict saddle point almost surely [\[Lee et al.,](#page-10-0) [2016,](#page-10-0) [Panageas and Piliouras,](#page-10-3) [2016\]](#page-10-3).

C Proof for Rank-1 Matrix Factorization (Theorem [3.2\)](#page-7-2)

In this section we prove Theorem [3.2.](#page-7-2)

Proof of Theorem [3.2.](#page-7-2) We define the following four key quantities:

$$
\alpha_t = \mathbf{u}_t^\top \mathbf{u}^*, \qquad \alpha_{t,\perp} = \|\mathbf{U}_{\perp}^* \mathbf{u}_t\|_2, \qquad \beta_t = \mathbf{v}_t^\top \mathbf{v}^*, \qquad \beta_{t,\perp} = \|\mathbf{V}_{\perp}^* \mathbf{v}_t\|_2,
$$

where $U_{\perp}^* = I - u^* u^{*\top}$ and $V_{\perp}^* = I - v^* v^{*\top}$ are the projection matrices onto the orthogonal complement spaces of u^* and v^* , respectively. Notice that $||u_t||_2^2 = \alpha_t^2 + \alpha_{t,\perp}^2$ and $||v_t||_2^2 =$ $\beta_t^2 + \beta_{t,\perp}^2$. It turns out that we can write down the explicit formulas for the dynamics of these quantities:

$$
\alpha_{t+1} = \left(1 - \eta \left(\beta_t^2 + \beta_{t,\perp}^2\right)\right) \alpha_t + \eta \sigma_1 \beta_t, \qquad \beta_{t+1} = \left(1 - \eta \left(\alpha_t^2 + \alpha_{t,\perp}^2\right)\right) \beta_t + \eta_1 \sigma_1 \alpha_t, \n\alpha_{t+1,\perp} = \left(1 - \eta \left(\beta_t^2 + \beta_{t,\perp}^2\right)\right) \alpha_{t,\perp}, \qquad \beta_{t+1,\perp} = \left(1 - \eta \left(\alpha_t^2 + \alpha_{t,\perp}^2\right)\right) \beta_{t,\perp}.
$$
\n(23)

To facilitate the analysis, we also define:

$$
h_t = \alpha_t \beta_t - \sigma_1,
$$

$$
\xi_t = \alpha_{t,\perp}^2 + \beta_{t,\perp}^2.
$$

Then our goal is to show $\xi_t \to 0$ and $h_t \to 0$ as $t \to \infty$. We calculate the dynamics of h_t and ξ_t :

$$
h_{t+1} = \left(1 - \eta \left(\alpha_t^2 + \beta_t^2\right) + \eta^2 \left(\alpha_t \beta_t h_t + \alpha_t^2 \beta_{t,\perp}^2 + \beta_t^2 \alpha_{t,\perp}^2 + \alpha_{t,\perp}^2 \beta_{t,\perp}^2\right)\right) h_t - \eta \alpha_t \beta_t \xi_t + \eta^2 \sigma_1 \alpha_{t,\perp}^2 \beta_{t,\perp}^2,
$$

$$
\xi_{t+1} = \left(1 - \eta \left(\beta_t^2 + \beta_{t,\perp}^2\right)\right)^2 \alpha_{t,\perp}^2 + \left(1 - \eta \left(\alpha_t^2 + \alpha_{t,\perp}^2\right)\right)^2 \beta_{t,\perp}^2.
$$
 (24)

According to our initialization scheme, with high probability we have $|\alpha_0|, |\beta_0| \in$ According to our initialization scheme, with high probability we have $|\alpha_0|, |\beta_0| \in$ 0.1 $c_{init}\sqrt{\frac{\sigma_1}{d}}$, 10 $c_{init}\sqrt{\frac{\sigma_1}{d}}$ and $|\alpha_{0,\perp}|, |\beta_{0,\perp}| \leq 10$ $c_{init}\sqrt{\sigma_1}$. We assume that these conditions are

satisfied. We also assume that the signal at the beginning is positive: $\alpha_0\beta_0 > 0$, which holds with probability 1/2. Without loss of generality we assume α_0 , $\beta_0 > 0$.^{[12](#page-19-0)}

We divide the dynamics into two stages.

Lemma C.1 (Stage 1: escaping from saddle point $(0, 0)$). *Let* $T_1 = \min \{ t \in \mathbb{N} : \alpha_t^2 + \beta_t^2 \geq \frac{1}{2}\sigma_1$ *. Then for* $t = 0, 1, \ldots, T_1 - 1$ *, the followings hold:*

- *(i) Positive signal strengths:* α_t , $\beta_t > 0$;
- (*ii*) *Small magnitudes in complement space:* $\xi_t \le \xi_0 \le 100c_{init}^2\sigma_1$; $\ddot{}$
- *(iii) Growth of magnitude in signal space:* $\left(1 + \frac{c_{step}}{3}\right)$ $(\alpha_t + \beta_t) \leq \alpha_{t+1} + \beta_{t+1} \leq$ $(1 + c_{step}) (\alpha_t + \beta_t);$
- *(iv) Bounded ratio between two layers:* $|\alpha_t \beta_t| \leq \frac{99}{101}(\alpha_t + \beta_t)$.

Furthermore, we have $T_1 = O(\log d)$ *.*

In this stage, the strengths in the complement spaces remain small $(\xi_t \leq \xi_0)$ and the strength in the signal space is growing exponentially $(\alpha_{t+1} + \beta_{t+1} \geq (1 + \frac{c_{step}}{3}) (\alpha_t + \beta_t))$. Furthermore, $|\alpha_t - \beta_t| \leq \frac{99}{101}(\alpha_t + \beta_t)$ implies $\frac{\alpha_t}{\beta_t} \in \left[\frac{1}{100}, 100\right]$, which means the signal strengths in the two layers are of the same order.

Then we enter stage 2, which is essentially a local convergence phase. The following lemma characterizes the behaviors of the strengths in the signal and noise spaces in this stage.

Lemma C.2 (Stage 2: convergence to global minimum). Let T_1 be as defined in Lemma [C.1.](#page-19-1) Then *there exists a universal constant* $c_1 > 0$ *such that the followings hold for all* $t \geq T_1$ *:*

- (a) *Non-vanishing signal strengths in both layers:* $\alpha_t, \beta_t \ge \sqrt{c_1 \sigma_1}$;
- *(b) Bounded signal strengths:* $\alpha_t \beta_t \leq \sigma_1$, *i.e.*, $h_t \leq 0$;
- *(c) Shrinking magnitudes in complement spaces:* $\xi_t \leq (1 c_1c_{step})^{t-T_1}\xi_0 \leq (1 c_1c_{step})^{t-T_1}$. $100c_{init}^2 \sigma_1$;
- *(d) Convergence in signal space:* $|h_{t+1}| \leq (1 c_1 c_{step})|h_t| + c_{step} \xi_t$.

Note that properties (a) and (b) in Lemma [C.2](#page-19-2) imply $c_0 \le \frac{\alpha_t}{\beta_t} \le C_0$ for all $t \ge T_1$, where c_0 , $C_0 > 0$ are universal constants. Property (c) implies that for all $t \ge T_1 + T_2$ where $T_2 = \Theta(\log \frac{1}{\epsilon})$, we have $\xi_t = O(\epsilon \sigma_1)$. Then property (d) tells us that after another $T_3 = O(\log \frac{1}{\epsilon})$ iterations, we can ensure $|h_t| = O(\epsilon \sigma_1)$ for all $t \ge T_1 + T_2 + T_3$. These imply $\left\| u_t v_t^\top - M^* \right\|_F = O(\epsilon \sigma_1)$ after $t = T_1 + T_2 + T_3 = O(\log \frac{d}{\epsilon})$ iterations, completing the proof of Theorem [3.2.](#page-7-2) \Box

Now we prove Lemmas [C.1](#page-19-1) and [C.2.](#page-19-2)

Proof of Lemma [C.1.](#page-19-1) We use induction to prove the following statements for $t = 0, 1, \ldots, T_1 - 1$:

$$
\mathcal{D}(t): \quad \alpha_t, \beta_t > 0,
$$
\n
$$
\mathcal{E}(t): \quad \xi_t \leq \xi_0 \leq 100 c_{init}^2 \sigma_1,
$$
\n
$$
\mathcal{F}(t): \quad \left(1 + \frac{c_{step}}{3}\right) (\alpha_t + \beta_t) \leq \alpha_{t+1} + \beta_{t+1} \leq (1 + c_{step}) (\alpha_t + \beta_t),
$$
\n
$$
\mathcal{G}(t): \quad |\alpha_t - \beta_t| \leq \frac{99}{101} (\alpha_t + \beta_t),
$$
\n
$$
\mathcal{H}(t): \quad \|\mathbf{u}_t\|^2 + \|\mathbf{v}_t\|^2 \leq \sigma_1.
$$

¹²If $\alpha_0, \beta_0 < 0$, we can simply flip the signs of u^* and v^* .

• Base cases.

We know that $\mathcal{D}(0), \mathcal{E}(0)$ and $\mathcal{G}(0)$ hold from our assumptions on the initialization.

• $\mathcal{D}(t), \mathcal{E}(t) \Longrightarrow \mathcal{F}(t) \quad (\forall t \leq T_1 - 1).$

From (23) we have

$$
\alpha_{t+1} + \beta_{t+1} = (1 + \eta \sigma_1) (\alpha_t + \beta_t) - \eta (\alpha_t^2 + \alpha_{t,\perp}^2) \beta_t - \eta (\beta_t^2 + \beta_{t,\perp}^2) \alpha_t
$$

\n
$$
\geq (1 + \eta \sigma_1 - \eta (\alpha_t^2 + \beta_t^2 + \xi_t)) (\alpha_t + \beta_t)
$$

\n
$$
\geq (1 + \eta \sigma_1 - \eta (\frac{\sigma_1}{2} + 100c_{init}^2 \sigma_1)) (\alpha_t + \beta_t)
$$

\n
$$
\geq (1 + \frac{\eta \sigma_1}{3}) (\alpha_t + \beta_t)
$$

\n
$$
= (1 + \frac{c_{step}}{3}) (\alpha_t + \beta_t),
$$

where in the second inequality we have used the definition of T_1 , and the last inequality is true when c_{init} is sufficiently small.

On the other hand we have

$$
\alpha_{t+1} + \beta_{t+1} = (1 + \eta \sigma_1) (\alpha_t + \beta_t) - \eta (\alpha_t^2 + \alpha_{t,\perp}^2) \beta_t - \eta (\beta_t^2 + \beta_{t,\perp}^2) \alpha_t
$$

\n
$$
\leq (1 + \eta \sigma_1) (\alpha_t + \beta_t)
$$

\n
$$
= (1 + c_{step}) (\alpha_t + \beta_t).
$$

• $\mathcal{E}(t) \Longrightarrow \mathcal{H}(t)$ ($\forall t \leq T_1 - 1$).

We have

$$
\|\mathbf{u}_t\|^2 + \|\mathbf{v}_t\|^2 = \alpha_t^2 + \beta_t^2 + \xi_t \le \frac{1}{2}\sigma_1 + 100c_{init}^2\sigma_1 \le \sigma_1.
$$

• $\mathcal{D}(t), \mathcal{H}(t) \Longrightarrow \mathcal{D}(t + 1) \quad (\forall t \leq T_1 - 1).$

From (23) we have

$$
\alpha_{t+1} = \left(1 - \eta \left\|\mathbf{v}_t\right\|^2\right) \alpha_t + \eta \sigma_1 \beta_t \geqslant \left(1 - \eta \sigma_1\right) \alpha_t = \left(1 - c_{step}\right) \alpha_t > 0.
$$

Similarly we have $\beta_{t+1} > 0$. Note that c_{step} is chosen to be sufficiently small.

• $\mathcal{H}(t) \Longrightarrow \mathcal{E}(t + 1) \quad (\forall t \leq T_1 - 1).$

Recall from [\(24\)](#page-18-2):

$$
\xi_{t+1} = \left(1 - \eta \left\|\bm{v}_t\right\|^2\right)^2 \alpha_{t,\perp}^2 + \left(1 - \eta \left\|\bm{u}_t\right\|^2\right)^2 \beta_{t,\perp}^2.
$$

Since $\eta \|v_t\|^2 \le \eta (\|u_t\|^2 + \|v_t\|^2) \le \eta \sigma_1 = c_{step} \le 1$ and $\eta \|u_t\|^2 \le 1$, we have $\xi_{t+1} \leq \alpha_{t,\perp}^2 + \beta_{t,\perp}^2 = \xi_t.$

• $\mathcal{D}(t), \mathcal{E}(t), \mathcal{F}(t), \mathcal{G}(t) \Longrightarrow \mathcal{G}(t + 1) \quad (\forall t \leq T_1 - 1).$ From [\(23\)](#page-18-1) we have

$$
\alpha_{t+1} - \beta_{t+1} = (1 - \eta \sigma_1)(\alpha_t - \beta_t) - \eta(\beta_t^2 + \beta_{t,\perp}^2)\alpha_t + \eta(\alpha_t^2 + \alpha_{t,\perp}^2)\beta_t = (1 - \eta \sigma_1 + \eta \alpha_t \beta_t)(\alpha_t - \beta_t) - \eta \beta_{t,\perp}^2 \alpha_t + \eta \alpha_{t,\perp}^2 \beta_t.
$$

From $\alpha_t^2 + \beta_t^2 < \frac{1}{2}\sigma_1$ we know $\alpha_t \beta_t < \frac{1}{4}\sigma_1$. Thus

$$
|\alpha_{t+1} - \beta_{t+1}| \leq (1 - \eta \sigma_1 + \eta \alpha_t \beta_t) |\alpha_t - \beta_t| + \eta \beta_{t,\perp}^2 \alpha_t + \eta \alpha_{t,\perp}^2 \beta_t
$$

$$
\leq (1 - \frac{3}{4} \eta \sigma_1) |\alpha_t - \beta_t| + \eta \xi_t (\alpha_t + \beta_t)
$$

$$
\leq \left(1 - \frac{3}{4}\eta\sigma_1\right) \cdot \frac{99}{101}(\alpha_t + \beta_t) + \eta \cdot 100c_{init}^2\sigma_1(\alpha_t + \beta_t) \n\leq \left(1 - \eta\sigma_1\left(\frac{3}{4} - 100c_{init}^2 \cdot \frac{101}{99}\right)\right) \cdot \frac{99}{101}(\alpha_t + \beta_t) \n\leq \frac{99}{101}(\alpha_t + \beta_t) \n\leq \frac{99}{101}(\alpha_{t+1} + \beta_{t+1}).
$$

 $\sqrt{2(\alpha_t^2 + \beta_t^2)} < \sqrt{2 \cdot \frac{1}{2} \sigma_1} =$ Lastly we upper bound T_1 . Note that for all $t < T_1$ we have $\alpha_t + \beta_t \leq$ $\sqrt{\sigma_1}$. From $\mathcal{F}(t)$ we know that $\alpha_t + \beta_t$ is increasing exponentially. Therefore, we must have $T_1 = O\left(\log \frac{\sqrt{\sigma_1}}{\alpha_0 + \beta}\right)$ $\overline{\frac{\sqrt{\sigma_1}}{\alpha_0 + \beta_0}}$ = $O\left(\log \frac{\sqrt{\sigma_1}}{\sqrt{\sigma_1/d}}\right)$ = $O(\log d)$. \Box

Proof of Lemma [C.2.](#page-19-2) By the definition of T_1 we know $\alpha_{T_1}^2 + \beta_{T_1}^2 \ge \frac{1}{2}\sigma_1$. In the proof of Lemma [C.1,](#page-19-1) we have shown $\alpha_{T_1}, \beta_{T_1} > 0$ and $|\alpha_{T_1} - \beta_{T_1}| \leq \frac{99}{101} (\alpha_{T_1} + \beta_{T_1})$. These imply $\min {\alpha_{T_1}, \beta_{T_1}} \geq$ we have shown $\alpha_{T_1}, \beta_{T_1} > 0$ and $|\alpha_{T_1} - \beta_{T_1}| \le 2\sqrt{c_1 \sigma_1}$ for some small universal constant $c_1 > 0$.

We use induction to prove the following statements for all $t \geq T_1$:

$$
\mathcal{I}(t): \qquad \alpha_t \geqslant \alpha_{T_1} \cdot \prod_{i=T_1}^{t-1} \left(1 - \eta \xi_0 \left(1 - c_1 c_{step}\right)^{i-T_1}\right), \quad \beta_t \geqslant \beta_{T_1} \cdot \prod_{i=T_1}^{t-1} \left(1 - \eta \xi_0 \left(1 - c_1 c_{step}\right)^{i-T_1}\right),
$$

$$
\mathcal{J}(t): \qquad \alpha_t, \beta_t \geqslant \sqrt{c_1 \sigma_1},
$$

$$
\mathcal{K}(t): \qquad \alpha_t \beta_t \leq \sigma_1, \text{ i.e., } h_t \leq 0,
$$

$$
\mathcal{L}(t): \qquad \xi_t \leq (1 - c_1 c_{step})^{t - T_1} \xi_0 \leq (1 - c_1 c_{step})^{t - T_1} \cdot 100 c_{init}^2 \sigma_1,
$$

 $\mathcal{M}(t) : \quad |h_{t+1}| \leq (1 - c_1 c_{step})|h_t| + c_{step} \xi_t.$

• Base cases.

 $\mathcal{I}(T_1)$ is obvious. We know that $\mathcal{J}(T_1)$ is true by the definition of c_1 . $\mathcal{K}(T_1)$ can be shown as follows:

$$
\alpha_{T_1} \beta_{T_1} \leq \frac{1}{4} \left(\alpha_{T_1} + \beta_{T_1} \right)^2
$$

\n
$$
\leq \frac{1}{4} \left(1 + c_{step} \right)^2 \left(\alpha_{T_1 - 1} + \beta_{T_1 - 1} \right)^2
$$
 (by Lemma C.1 (iii))
\n
$$
\leq \frac{1}{4} \left(1 + c_{step} \right)^2 \cdot 2 \left(\alpha_{T_1 - 1}^2 + \beta_{T_1 - 1}^2 \right)
$$

\n
$$
\leq \frac{1}{4} \left(1 + c_{step} \right)^2 \cdot 2 \cdot \frac{1}{2} \sigma_1
$$
 (by the definition of T_1)
\n
$$
\leq \sigma_1.
$$
 (choosing c_{step} to be small)

 $\mathcal{L}(T_1)$ reduces to $\xi_{T_1} \leq \xi_0$, which was shown in the proof of Lemma [C.1.](#page-19-1)

• $\mathcal{I}(t) \Longrightarrow \mathcal{J}(t)$ ($\forall t \geq T_1$).

Notice that we have $\eta \xi_0 \leq \frac{c_{step}}{\sigma_1}$ $\frac{f_{step}}{\sigma_1} \cdot 100c_{init}^2 \sigma_1 = 100c_{step}c_{init}^2 < \frac{1}{2}$ since c_{step} and c_{init} are sufficiently small. Then we have

$$
\alpha_{t} \geq \alpha_{T_{1}} \cdot \prod_{i=T_{1}}^{t-1} \left(1 - \eta \xi_{0} \left(1 - c_{1} c_{step} \right)^{i-T_{1}} \right)
$$

\n
$$
\geq \alpha_{T_{1}} \cdot \prod_{i=0}^{\infty} \left(1 - \eta \xi_{0} \left(1 - c_{1} c_{step} \right)^{i} \right)
$$

\n
$$
\geq \alpha_{T_{1}} \cdot \prod_{i=0}^{\infty} \exp \left(-2\eta \xi_{0} \left(1 - c_{1} c_{step} \right)^{i} \right) \qquad (1 - x \geq e^{-2x}, \forall 0 \leq x \leq 1/2)
$$

$$
= \alpha_{T_1} \cdot \exp\left(-\frac{2\eta \xi_0}{c_1 c_{step}}\right)
$$

\n
$$
\geq \alpha_{T_1} \cdot \exp\left(-\frac{200 c_{step} c_{init}^2}{c_1 c_{step}}\right)
$$

\n
$$
\geq 2\sqrt{c_1 \sigma_1} \cdot \exp\left(-\frac{200 c_{init}^2}{c_1}\right)
$$

\n
$$
\geq \sqrt{c_1 \sigma_1}.
$$

(choosing c_{init} to be small)

Similarly we have $\beta_t \geqslant \sqrt{c_1 \sigma_1}$.

• $\mathcal{I}(t), \mathcal{J}(t), \mathcal{K}(t), \mathcal{L}(t) \Longrightarrow \mathcal{I}(t + 1) \quad (\forall t \geq T_1).$ From (23) we have $\alpha_{t+1} = \left(1 - \eta\right)$ $\beta_t^2 + \beta_{t,\perp}^2$ $\alpha_t + \eta \sigma_1 \beta_t$ $=$ $\overline{}$ $1 - \eta \beta_{t,\perp}^2$ $\alpha_t - \eta h_t \beta_t$ $\overline{}$ \overline{y}

$$
\geq (1 - \eta \beta_{t, \perp}^2) \alpha_t
$$

\n
$$
\geq (1 - \eta \xi_t) \alpha_t
$$

\n
$$
(h_t \leq 0, \beta_t > 0)
$$

$$
\geq (1 - \eta \xi_0 (1 - c_1 c_{step})^{t - T_1}) \alpha_t \tag{L(t)}
$$

$$
\geqslant \alpha_{T_1} \cdot \prod_{i=T_1}^{r} \left(1 - \eta \xi_0 \left(1 - c_1 c_{step}\right)^{i-T_1}\right). \tag{I(t)}
$$

Similarly we have $\beta_{t+1} \geq \beta_{T_1} \cdot \prod_{i=1}^{t}$ $i = T_1$ $1 - \eta \xi_0 \left(1 - c_1 c_{step} \right)^{i - T_1}$.

• $\mathcal{J}(t), \mathcal{K}(t), \mathcal{L}(t) \Longrightarrow \mathcal{K}(t + 1) \ (\forall t \geq T_1).$

From (24) we have

$$
h_{t+1} = \left(1 - \eta \left(\alpha_t^2 + \beta_t^2\right) + \eta^2 \left(\alpha_t \beta_t h_t + \alpha_t^2 \beta_{t,\perp}^2 + \beta_t^2 \alpha_{t,\perp}^2 + \alpha_{t,\perp}^2 \beta_{t,\perp}^2\right)\right) h_t - \eta \alpha_t \beta_t \xi_t + \eta^2 \sigma_1 \alpha_{t,\perp}^2 \beta_{t,\perp}^2
$$

\$\leq \left(1 - \eta \left(\alpha_t^2 + \beta_t^2\right)\right) h_t + \eta^2 \alpha_t \beta_t h_t^2 - \eta \alpha_t \beta_t \xi_t + \eta^2 \sigma_1 \alpha_{t,\perp}^2 \beta_{t,\perp}^2, (25)

where we have used $h_t \le 0$. Since $\alpha_t, \beta_t \ge \sqrt{c_1 \sigma_1}$ and $\alpha_t \beta_t \le \sigma_1$, we have $\alpha_t, \beta_t \in C$. $\Theta(\sqrt{\sigma_1})$. Furthermore, we can choose c_{step} and c_{init} small enough such that $\eta \xi_0 \leq 4c_1$ which implies

$$
\eta^2 \sigma_1 \alpha_{t,\perp}^2 \beta_{t,\perp}^2 \leq \eta^2 \sigma_1 \cdot \frac{1}{4} \xi_t^2 \leq \frac{1}{4} \eta \xi_t \cdot \eta \sigma_1 \xi_0 \leq \eta \xi_t \cdot c_1 \sigma_1 \leq \eta \xi_t \cdot \alpha_t \beta_t.
$$

Therefore [\(25\)](#page-22-0) implies

$$
h_{t+1} \leq (1 - \eta \cdot O(\sigma_1)) h_t + \eta^2 \alpha_t \beta_t h_t^2
$$

=
$$
(1 - \eta \cdot O(\sigma_1) + \eta^2 \alpha_t \beta_t h_t) h_t
$$

$$
\leq (1 - \eta \cdot O(\sigma_1) - \eta^2 \sigma_1^2) h_t \qquad (0 < \alpha_t \beta_t \leq \sigma_1)
$$

=
$$
(1 - O(c_{step}) - c_{step}^2) h_t
$$

$$
\leq 0,
$$

where the last step is true when c_{step} is sufficiently small.

• $\mathcal{J}(t), \mathcal{K}(t), \mathcal{L}(t) \Longrightarrow \mathcal{L}(t + 1) \quad (\forall t \geq T_1).$

From $\alpha_t, \beta_t \ge \sqrt{c_1 \sigma_1}$ and $\alpha_t \beta_t \le \sigma_1$ we have $\alpha_t, \beta_t = \Theta(\sqrt{\sigma_1})$. Also we have $\xi_t \le \xi_0$. Thus we can make sure $\eta(\alpha_t^2 + \alpha_{t,\perp}^2) < 1$ and $\eta(\beta_t^2 + \beta_{t,\perp}^2) < 1$. Then from [\(24\)](#page-18-2) we have

$$
\xi_{t+1} = \left(1 - \eta \left(\beta_t^2 + \beta_{t,\perp}^2\right)\right)^2 \alpha_{t,\perp}^2 + \left(1 - \eta \left(\alpha_t^2 + \alpha_{t,\perp}^2\right)\right)^2 \beta_{t,\perp}^2
$$

\n
$$
\leq \left(1 - \eta \beta_t^2\right)^2 \alpha_{t,\perp}^2 + \left(1 - \eta \alpha_t^2\right) \beta_{t,\perp}^2
$$

\n
$$
\leq \left(1 - \eta c_1 \sigma_1\right) \xi_t
$$

\n
$$
= \left(1 - c_1 c_{step}\right) \xi_t.
$$

• We have shown $\mathcal{I}(t),\mathcal{J}(t),\mathcal{K}(t)$ and $\mathcal{L}(t)$ for all $t \geq T_1$. Now we use them to prove $\mathcal{M}(t)$ for all $t \geqslant T_1$:

$$
|h_{t+1}| = (1 - \eta \left(\alpha_t^2 + \beta_t^2\right) + \eta^2 \left(\alpha_t \beta_t h_t + \alpha_t^2 \beta_{t,\perp}^2 + \beta_t^2 \alpha_{t,\perp}^2 + \alpha_{t,\perp}^2 \beta_{t,\perp}^2\right)) |h_t| + \eta \alpha_t \beta_t \xi_t - \eta^2 \sigma_1 \alpha_{t,\perp}^2 \beta_{t,\perp}^2
$$

\$\leq \left(1 - \frac{1}{2} \eta \left(\alpha_t^2 + \beta_t^2\right)\right) |h_t| + \eta \alpha_t \beta_t \xi_t\$
\$\leq \left(1 - \frac{1}{2} \eta \cdot 2c_1 \sigma_1\right) |h_t| + \eta \sigma_1 \xi_t\$
= (1 - c_1 c_{step}) |h_t| + c_{step} \xi_t.

Here we have used $\eta \leq \frac{\alpha_t^2 + \beta_t^2}{2|\alpha_t \beta_t h_t + \alpha_t^2 \beta_{t,1}^2 + \beta_t^2 \alpha_{t,1}^2 + \alpha_{t,1}^2 \beta_{t,1}^2|}$, which is clearly true when c_{step} is small enough.

Therefore, we have finished the proof of Lemma [C.2.](#page-19-2)

 \Box